

**SCIENCO**  
**SOUTHERN BRAZILIAN JOURNAL**  
**OF CHEMISTRY**  
ISSN 0104-5431

**AN INTERNATIONAL FORUM FOR THE RAPID PUBLICATION OF ORIGINAL  
SCIENTIFIC ARTICLES DEALING WITH CHEMISTRY AND RELATED  
INTERDISCIPLINARY AREAS**

**VOLUME TWENTY-FOUR, NUMBER TWENTY-FOUR**

**DECEMBER 2015**

## EDITOR

LAVINEL G. IONESCU, *SCIENCO*, Consultoria Científica, Viamão, RS, BRASIL

## ASSISTANT EDITOR

LUIS ALCIDES BRANDINI DE BONI, Tchequímica LTDA, Porto Alegre, RS, Brasil.

## EDITORIAL BOARD

- FARUK NOME AGUILERA, Departamento de Química, Universidade Federal de Santa Catarina, Florianópolis, SC, Brasil
- D. BALASUBRAMANIAN, Centre for Cellular and Molecular Biology, Hyderabad, INDIA
- HECTOR E. BERTORELLO, Departamento de Química Orgánica, Facultad de Ciencias Químicas, Universidad Nacional de Córdoba, Córdoba, ARGENTINA
- AÉCIO P. CHAGAS, Instituto de Química, UNICAMP, Campinas, SP, BRASIL
- JUAN JOSÉ COSA, Departamento de Química y Física, Facultad de Ciencias Exactas, Universidad Nacional de Río Cuarto, Río Cuarto, ARGENTINA
- GLENN A. CROSBY, Department of Chemistry, Washington State University, Pullman, WA, USA
- VITTORIO DEGIORGIO, Dipartimento di Elettronica, Sezione di Fisica Applicata, Università di Pavia, Pavia, ITALIA
- JOSE C. TEIXEIRA DIAS, Departamento de Química, Universidade de Coimbra, Coimbra, PORTUGAL
- OMAR A. EL SEOUD, Instituto de Química, Universidade de São Paulo, São Paulo, SP, BRASIL
- FERNANDO GALEMBECK, Instituto de Química, UNICAMP, Campinas, SP, BRASIL
- NISSIM GARTI, Casali Institute of Applied Science, Hebrew University of Jerusalem, Jerusalem, ISRAEL
- GASPAR GONZALEZ, Centro de Pesquisa, CENPES-PETROBRAS, Ilha do Fundão, Rio de Janeiro, RJ, BRASIL
- YOSHITAKA GUSHIKEM, Instituto de Química, UNICAMP, Campinas, SP, BRASIL
- WILLIAM HASE, Department of Chemistry, Texas Tech University, Lubbock, Texas, USA
- I. B. IVANOV, Laboratory of Thermodynamics and Physico-chemical Hydrodynamics, Faculty of Chemistry, University of Sofia, Sofia, BULGARIA
- IVAN IZQUIERDO, Departamento de Bioquímica, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, BRASIL
- V.A. KAMINSKY, Karpov Institute of Physical Chemistry, Moscow, RUSSIA
- MICHAEL LAING, Department of Chemistry, University of Natal, Durban, SOUTH AFRICA
- EDUARDO LISSI, Departamento de Química, Universidad de Santiago de Chile, Santiago, CHILE
- WALTER LWOWSKI, Department of Chemistry, New Mexico State University, Las Cruces, N.M., USA
- CRISTINA MANDRAVEL, Catedra de Chimie Fizica, Facultatea de Chimie Universitatea din Bucuresti, Bucuresti, Romania
- C. MANOHAR, Bhabha Atomic Research Centre, Chemistry Division, Bombay, INDIA
- AYRTON FIGUEIREDO MARTINS, Departamento de Química, Universidade Federal de Santa Maria, Santa Maria, RS, BRASIL
- FRED MENDER, Department of Chemistry, Emory University, Atlanta, GA, USA
- KASHMIRI LAL MITTAL, Private Consultant, Hopewell Junction, N.Y., USA
- ARNO MULLER, Escola de Engenharia, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, BRASIL
- JOSE MIGUEL PARERA, Instituto de Investigaciones en Catalisis y Petroquímica, Universidad Nacional del Litoral, Santa Fe, ARGENTINA
- LARRY ROMSTED, Department of Chemistry, Rutgers University, Piscataway N.J., USA
- GILBERTO FERNANDES DE SA, Departamento de Química Fundamental, Universidade Federal de Pernambuco, Recife, PE, BRASIL
- DIMITRIOS SAMIOS, Instituto de Química, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, BRASIL
- DIOGENES DOS SANTOS, Department of Molecular Biology, Oxford University, Oxford, ENGLAND
- BEN K. SELINGER, Department of Chemistry, Australian National University, Canberra, AUSTRALIA
- KOZO SHINODA, Department of Applied Chemistry, Faculty of Engineering, Yokohama National University, Yokohama, JAPAN
- CRISTOFOR I. SIMIONESCU, Academia Romana, Filiala Iasi, Iasi, ROMANIA
- BRUNO SZPOGANICZ, Departamento de Química, Universidade Federal de Santa Catarina, Florianópolis, SC, Brasil
- UMBERTO TONELLATO, Dipartimento di Chimica Organica, Università degli Studi di Padova, Padova, ITALIA
- DIETER VOLLHARDT, Max Planck Institut für Kolloid und Grenzflächenforschung, Berlin, GERMANY
- RAOUL ZANA, Institut Charles Sadron, CRM-EAHP, Strassbourg, FRANCE

## INFORMATION FOR AUTHORS

The Southern Brazilian Journal of Chemistry - SBJC will publish review articles, original research papers and short communications dealing with chemistry and interdisciplinary areas such as materials science, biotechnology, bioengineering and other multidisciplinary fields.

Articles report the results of a complete study. They should include an Abstract, Introduction describing the known art in the field Experimental or Materials and Methods, Results and Discussion, Acknowledgments (when appropriate) and References.

Short Communications should be limited to 1500 words, including the equivalent space for figures and/or tables and should include an Abstract and concise Experimental.

Manuscripts may be submitted on-line or in triplicate (original and two copies by registered mail) and are received with the understanding that the original has not been submitted for publication elsewhere. It is implicit that all the persons listed as authors have given their approval for the submission of the paper and that permission has also been granted by the employer, when necessary.

Manuscripts must be written in American or British English, single spaced, on A4 sheets (21 cm x 29.5 cm) and one side only and should be numbered beginning with the title page. Type must be 12 Arial or Times New Roman.

Margins of at least 3 cm should be left at the top and bottom and both sides of each page. The first page of the paper should contain only the title of the paper, the name(s) and addressees) of the author(s), an abstract of not more than 250 words and 4-8 keywords. We reserve the right to translate the abstract in Portuguese. Abstracts are required of all papers including reviews and short communications.

Figures and Tables with short explanatory titles, each on a separate sheet, should be adequate for direct reproduction and identified in pencil on the back of each page by Arabic numerals, corresponding to the order they appear in the manuscript. Tables and Figures (BMP or JPG format) may also be included directly in the text when convenient and the article may be submitted in a quasi-final form in order to facilitate editorial work.

References should be numbered in the sequence they appear in the text, cited by superior numbers and listed at the end of the paper in the reference section in the numerical order they appear in the text. The style for references is shown below:

1. L. G. Ionescu and D. S. Fung, *J. Chem. Soc. Faraday Trans. 1*, 77, 2907-2912 (1981).
2. K. L. Mittal, Ed., "*Solution Chemistry of Surfactants*", Plenum Press, New York (1984), Vols. 1-3, pp. 1-2173.

IUPAC Rules should be used for the name of chemical compounds and preference should be given to 51 units.

Authors are invited to send manuscripts by registered air mail to the EDITOR - SBJC, C.P. 15032, Agronomia, Porto Alegre, RS BRASIL 91501, or by e-mail to

[lavinel@gmail.com](mailto:lavinel@gmail.com)

[www.sbjchem.tk](http://www.sbjchem.tk)

# SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY

ISSN 0104-543

VOLUME TWENTY THREE, NUMBER TWENTY THREE

2015

## CONTENTS / CONTEÚDO

EDITORIAL .....	1
CARACTERIZAÇÃO HIDROGEOQUÍMICA DOS ÍONS PREDOMINANTES NAS ÁGUAS SUBTERRÂNEAS DA MINA 3 (COMPANHIA NITROQUÍMICA BRASILEIRA), MORRO DA FUMAÇA, SANTA CATARINA, BRASIL C. Kath, P. C. P. das Neves and L. G. Ionescu.....	2
VOLATILE COMPONENTS OF <i>ONCOBA SPINOSA</i> LEAF AND <i>MORUS MESOZYGIA</i> LEAF AND STEM BARKS. A. Aboaba, O. Adeeko and G. Flamini .....	23
THE CHEMICAL AND MEDICINAL POTENTIALS OF THE FRUIT ESSENTIAL OIL OF <i>Chrysophyllum cainito</i> (INDIA STAR APPLE). O. Ogumola, O. Bolaji, O. S. & Sodame, A. ....	37
SYNTHESIS AND BIOLOGICAL ACTIVITIES EVALUATION OF SOME NEW SPIRO 1,2,4-TRIAZOLE DERIVATIVES HAVING SULFONAMIDE MOIETY. Hany M. Dalloul, Khaled El-nwairy, b Ali Z. Shoraf, Ahmed Abu Samaha.....	49
KINETICS OF REACTION BETWEEN MALACHITE GREEN AND HYDROXYL ION IN THE PRESENCE OF REDUCING SUGARS. Latona, D. Felix and Dada, A. Oluwasogo .....	61
CHEMICAL CONSTITUENTS AND PHYTOCHEMICAL STUDIES OF EPAZOTE ( <i>Chenopodium ambrosioides</i> ) LINN GROWN IN NORTH CENTRAL NIGERIA Larayetán, R. A., Adegbe A. A., Owonikoko A. D. ....	73
VOLATILE OIL CONSTITUENTS OF THE LEAVES AND WOODS OF <i>Cedrela</i> <i>odorata</i> AND <i>Dalbergia latifolia</i> FROM SOUTH WEST NIGERIA. Ugah, T. C, Aboaba, S. A and Ekundayo, O .....	83
CHEMICAL COMPOSITIONS AND ANTIMICROBIAL ACTIVITY OF THE LEAF ESSENTIAL OIL OF <i>Gossypium hirsutum</i> Ogunmola Oluranti O. ....	91
BOOKS / LIVROS .....	101
AUTHOR INDEX .....	109

## **EDITORIAL**

With this issue, THE SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY completes twenty three (23) years of continuous, uninterrupted publication. The printed version of the JOURNAL has been distributed in more than fifty (50) countries of the globe, covering all continents.

We publicly thank all the members of the Editorial Board that includes distinguished scientists from all continents for the assistance, advice and financial help given during the last twenty three (23) years.

At the present, because of poor health we can not continue as Editor.

Professor Dr. Luis Brandini De Boni is assuming as Editor in Chief of the SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY.

Prof. Dr. Lavinel G. Ionescu.

**CARACTERIZAÇÃO HIDROGEOQUÍMICA DOS ÍONS PREDOMINANTES  
NAS ÁGUAS SUBTERRÂNEAS DA MINA 3 (COMPANHIA NITROQUÍMICA  
BRASILEIRA), MORRO DA FUMAÇA, SANTA CATARINA, BRASIL**

**Cibele Kath; Paulo César Pereira das Neves – Laboratório de Geologia e  
Mineralogia, Curso de Química Industrial  
Universidade Luterana do Brasil, Canoas, RS, Brasil  
usppd@yahoo.com.br**

**Lavinel G. Ionescu  
SCIENCO Scientific Consulting Services, Viamão, RS, Brasil  
and  
Sarmisegetuza Research Group, Santa Fe, New Mexico, USA  
Lavinel.g.ionescu@gmail.com**

**ABSTRACT**

**This paper presents a comparative hydrogeochemical characterization on the predominant ions of the underground water that flow the Cocal vein, in the Mina 3 (Companhia Nitroquímica Brasileira), Fluorine District of Santa Catarina, Morro da Fumaça Co., Santa Catarina, Brazil. Four samples were selected: P<sub>1</sub> – underground water to 300 m of depth; P<sub>2</sub> – underground water to 150 m of depth; P<sub>3</sub> – water of the dam of reject; P<sub>4</sub> – treated water of provisioning of Companhia Catarinense de Águas e Saneamento (CASAN). In the water sample collected in the selected sources, quantitative determinations of physical-chemical parameters (pH and electric conductivity) and of cations (Al, Ca, Cl, Li, Mg, Mn, K, Si, Na, and Zn) and anions (sulfate, chloride, bicarbonate, and fluorine) were performed. The results obtained revealed that most of the analyzed ions are related to the representatives originated from rocks that constitute the pools where the studied water flow. The hydrogeochemical classification was obtained starting from the milliequivalence pattern among the main elements, in the cations and anions form,**

C. Kath, P. C. P. das Neves and L. G. Ionescu

being used the triangular diagram of Piper. The underground water of the point 1 presented a hydrogeochemical characterization compatible with its geological environment occurrence and being perceptible the characterization of an interrelation among the rock and the water that circulate within and they also presented an amount of larger ions than the water of the point 2, in since it flow through a mineralized pool. The underground water of the point 2 presented good potable conditions, with some restrictions. In the water of the point 3 there is a larger ions concentration by due to the washing of the mineral, that allows a larger solubilization of the elements, and the time of permanence of the water in the lake also causes large concentration. The water collected in P<sub>4</sub> point exhibit good water potability parameters and absence of conditions that favor fluorosis.

## RESUMO

O artigo apresenta a caracterização hidrogeoquímica dos íons predominantes nas águas de percolação do filão Cocal, Mina 3 (Companhia Nitroquímica Catarinense), Distrito Fluorítico de Santa Catarina, Morro da Fumaça, Santa Catarina, Brasil. Foram selecionados quatro pontos de amostragem: P<sub>1</sub> – águas subterrâneas a 300 m de profundidade; P<sub>2</sub> – águas subterrâneas a 150 m de profundidade; P<sub>3</sub> – águas da barragem de rejeito; P<sub>4</sub> – água de abastecimento da mina, tratada pela Companhia Catarinense de Águas e Saneamento (CASAN). Nas amostras coletadas nos pontos selecionados foram realizadas determinações quantitativas de parâmetros físico-químicos (pH e condutividade elétrica) e de cátions e ânions predominantes (Al, Ca, Cl, Li, Mg, Mn, K, Si, Na, Zn, sulfato, cloreto, bicarbonato e fluoreto). Os resultados obtidos revelaram que a maioria dos íons analisados apresentam relação direta com os constituintes advindos das rochas que constituem os aquíferos por onde as águas circulam. A classificação hidrogeoquímica foi obtida a partir da relação de miliequivalência entre os elementos principais, na forma de cátions e ânions, utilizando-se o diagrama triangular de Piper. As águas P<sub>1</sub> apresentaram

caracterização hidrogeoquímica compatível com seu ambiente de ocorrência, sendo perceptível a caracterização de uma interrelação entre as rochas e as águas que nelas circula; nas águas P2 a quantidade de íons foi menor do que em P1, pelo fato de circularem em um aquífero mais mineralizado; as águas P<sub>3</sub> apresentam uma concentração de íons maior, fator que se deve à lavagem do minério, com conseqüente maior solubilização dos elementos químicos, além do que, uma maior concentração é devido ao maior tempo de permanência das mesmas no lago; as águas P4 apresentam bons parâmetros de potabilidade, além de ausência de condições que favoreçam fluorose.

## INTRODUCTION

Os organismos podem privar-se de água de alimentação por poucos dias, entretanto privando-os de água os mesmos podem sucumbir em poucas horas. Nas últimas décadas têm-se notado uma crescente preocupação em relação à escassez dos recursos hídricos. Em áreas industrializadas encontra-se uma forte marca das atividades humanas na qualidade química das águas. Este fato é em particular marcante onde predominam aquíferos passíveis de serem influenciados pelas atividades antrópicas. Nas proximidades dos grandes centros urbanos temos, principalmente, problemas associados às descargas de poluentes. Em áreas agrícolas, a química da água pode estar fortemente influenciada pelos produtos químicos utilizados e nas regiões de mineração temos um caso especial de indústria que acarreta um impacto significativo sobre a qualidade das águas subterrâneas. O homem dispõe naturalmente de dois contingentes de águas disponíveis para uso imediato: as águas de escoamento superficial, disponíveis em rios e lagos (cerca de 3%) e as águas subterrâneas (cerca de 97%)<sup>(1,2)</sup>. As características das águas subterrâneas refletem os meios por onde elas percolam, guardando uma estreita relação com os tipos de rochas drenadas e com os produtos das atividades humanas, adquiridos ao longo de seu trajeto. Todas as águas na natureza possuem, em graus distintos, um conjunto de sais em solução, sendo que as águas subterrâneas possuem, em geral, teores mais elevados do que as águas de superfície, por estarem intimamente expostas aos materiais solúveis presentes nos

C. Kath, P. C. P. das Neves and L. G. Ionescu

solos e nas rochas. A quantidade e o tipo de sais presentes nestas águas dependem, em consequência, do meio percolado, do tipo e velocidade do fluxo subterrâneo, da fonte de recarga do aquífero e do clima vigente na região.

Este trabalho apresenta os resultados obtidos em análises feitas em amostras de água provenientes de quatro pontos de referência da Mina 3 (Companhia Nitroquímica Catarinense), Morro da Fumaça Co., Santa Catarina, Brasil.

As amostras P<sub>1</sub> e P<sub>2</sub> foram coletadas respectivamente a 300 e 150 m de profundidade, nas galerias da mina, que percolam respectivamente o filão de fluorita Cocal e as coberturas sedimentares sobrejacentes ao Granito Pedras Grandes; P<sub>3</sub> - água do lago de decantação (barragem de rejeitos) da mineradora; P<sub>4</sub> - água de abastecimento da mineradora, tratada pela Companhia Catarinense de Águas e Saneamento (CASAN). Foram analisados dez cátions (alumínio, cálcio, ferro, lítio, magnésio, manganês, potássio, silício, sódio e zinco) e quatro ânions (bicarbonato, cloreto, fluoreto e sulfato), além do pH e da condutividade elétrica dessas águas.

Os objetivos do trabalho foram: caracterizar as águas provenientes da Mina 3 quanto aos aspectos físico-químicos e quanto aos íons predominantes; classificar hidrogeoquimicamente as águas dos quatro pontos considerados; verificar se a concentração dos parâmetros analisados nas águas para consumo humano atendem aqueles estabelecidos pela resolução do CONAMA (n<sup>o</sup> 396, 03.04.2008)<sup>(3)</sup>; comparar a potabilidade da água que abastece o município de Morro da Fumaça com as águas subterrâneas servidas na Vila Mina Fluorita (moradores da mineradora).

## LOCAL ESTUDADO

O presente estudo foi desenvolvido no local geologicamente identificado como filão Cocal (28<sup>o</sup>36'42''S; 49<sup>o</sup>15'58''W), Distrito Fluorítico de Santa Catarina, Bacia do rio Urussanga, Folha de Criciúma (SH.22-X-B-IV-1), Companhia

Nitroquímica Catarinense (Grupo Votorantim), Morro da Fumaça, Santa Catarina, Brasil.

## GEOLOGIA

O embasamento do Distrito Fluorítico da Santa Catarina é constituído predominantemente pelo Granito Pedras Grande<sup>(4)</sup>, que localmente é cortado por diques de rochas subvulcânicas ácidas do Eo-Paleozóico<sup>(5)</sup>. Também ocorrem rochas da Bacia do Paraná (formações Rio do Sul e Rio Bonito – arenitos e lamitos), além de diques e soleiras de diabásio da Formação Serra Geral. Todas estas litologias são cortadas pelo filão Cocal (fluorita), que se encaixa preferencialmente no granito (Figure 1).




Figure 1 – Geological map of the study location (modified)<sup>(6)</sup>.

C. Kath, P. C. P. das Neves and L. G. Ionescu

Os principais minerais ocorrentes ao longo da Mina 3 são: fluorite ( $\text{CaF}_2$ ), illite  $((\text{K},\text{H}_3\text{O})(\text{Al},\text{Mg},\text{Fe})_2(\text{Si},\text{Al})_4\text{O}_{10}[(\text{OH})_2,\text{H}_2\text{O}])$ -montmorillonite (smectite)  $(\text{Na},\text{Ca})_{0,3}(\text{Al},\text{Mg})_2\text{Si}_4\text{O}_{10}(\text{OH})_2.n\text{H}_2\text{O}$ , barite ( $\text{BaSO}_4$ ), quartz ( $\text{SiO}_2$ ), pyrite ( $\text{FeS}_2$ ), goethite ( $\alpha\text{-FeO}(\text{OH})$ ), hematita ( $\text{Fe}_2\text{O}_3$ ) e psilomelane (a mixture of manganese oxides)<sup>(6-7)</sup>.

## MATERIAIS E MÉTODOS

A etapa de campo constou de uma visita à Mina 3 (Companhia Nitroquímica Brasileira), no município de Morro da Fumaça, para reconhecimento da mineradora e coleta das amostras de água.

As amostras foram coletadas em garcos de polietileno com tampa, com capacidade de 1 L cada. Depois de coletadas, as mesmas foram armazenadas em ambiente de refrigeração até o instante da realização dos procedimentos analíticos, que seguiram a preconização do Standad Methods<sup>(8)</sup>. As amostras foram subdivididas em volumes menores, sendo que as destinadas às determinações de cátions foram preservadas em ácido nítrico concentrado. Para as determinações dos parâmetros físico-químicos e as análises de cátions e ânions, foram utilizados os seguintes equipamentos: estufas, sistema ultrapurificador de água, forno de mufla, balanças analíticas, pHmetro de bancada, condutivímetro, sensor de temperatura e eletrodo seletivo para fluoreto, chapas de agitação, espectômetro de absorção atômica, forno de grafite e fotômetro de chama. Para eliminar possíveis contaminantes iônicos a vidraria foi previamente lavada e imersa em solução de ácido nítrico  $0.1 \text{ mol.L}^{-1}$ .

As amostras sofreram digestões ácidas (branda (para metais solúveis) e enérgica (para metais totais)) com o intuito de disponibilizar em matrizes solúveis os elementos metálicos constituintes. Após, seguiu-se os procedimentos analíticos com as diversas determinações paramétricas que visaram avaliar o comportamento de cada elemento analisado e suas interações com os demais elementos em relação ao meio percolado por essas águas.

## RESULTADOS

Os resultados obtidos nos procedimentos analíticos estão apresentados nas tabelas abaixo:

Tabela 1 – Parâmetros físico-químicos

	Point 1	Point 2	Point 3	Point 4
pH	7.35	6.53	7.33	6.50
Condutividade elétrica ( $\mu\text{S}\cdot\text{cm}^{-1}$ )	500	200	516	712

Tabela 2 – Teor de ânions ( $\text{mg}\cdot\text{L}^{-1}$ )

	Point 1	Point 2	Point 3	Point 4
Dicarbonate	99.30	52.30	131.40	19.00
Clorine	69.42	23.14	54.00	15.42
Fluorine	7.31	1.67	14.29	0.42
Sulfate	0.36	0.25	0.45	0.14

Tabela 3 – Teor de cátions disponíveis ( $\text{mg}\cdot\text{L}^{-1}$ )

	Point 1	Point 2	Point 3	Point 4
Aluminum	0.307 $\pm$ 0.012	0.207 $\pm$ 0.006	7.053 $\pm$ 0.006	0.273 $\pm$ 0.006
Calcium	30.467 $\pm$ 0.231	10.882 $\pm$ 0.017	26.900 $\pm$ 0.100	2.709 $\pm$ 0.002
Iron	0.540 $\pm$ 0.005	0.275 $\pm$ 0.004	17.160 $\pm$ 0.010	0.672 $\pm$ 0.004
Lítio	0.108 $\pm$ 0.002	0.039 $\pm$ 0.00	30.108 $\pm$ 0.002	0.019 $\pm$ 0.002
Magnesium	1.893 $\pm$ 0.006	2.303 $\pm$ 0.006	3.947 $\pm$ 0.006	1.917 $\pm$ 0.006
Manganese	0.045 $\pm$ 0.001	0.027 $\pm$ 0.001	0.581 $\pm$ 0.001	0.027 $\pm$ 0.001
Potassium	2.090 $\pm$ 0.000	1.184 $\pm$ 0.000	7.321 $\pm$ 0.000	1.285 $\pm$ 0.000
Silicon	18.041 $\pm$ 0.165	18.356 $\pm$ 0.098	29.686 $\pm$ 0.197	5.603 $\pm$ 0.204
Sodium	46.516 $\pm$ 0.000	15.206 $\pm$ 0.000	61.308 $\pm$ 0.000	9.585 $\pm$ 0.000
Zinc	0.088 $\pm$ 0.00	10.314 $\pm$ 0.010	0.563 $\pm$ 0.001	10.974 $\pm$ 0.006

C. Kath, P. C. P. das Neves and L. G. Ionescu

Tabela 4 - Teor de cátions disponíveis (mg.L<sup>-1</sup>)

	Point 1	Point 2	Point 3	Point 4
Aluminum	0.537±0.015	0.397±0.006	20.757±0.012	0.390±0.000
Calcium	63.167±0.076	13.032±0.010	36.100±0.000	3.117±0.002
Iron	1.094±0.002	1.163±0.004	26.400±0.017	0.934±0.003
Lítio	0.130±0.002	0.050±0.000	0.228±0.000	0.040±0.003
Magnesium	2.060±0.000	2.787±0.006	5.290±0.000	2.940±0.000
Manganese	0.051±0.001	0.062±0.002	0.769±0.001	0.062±0.002
Potassium	10.703±0.065	1.687±0.000	16.677±0.000	1.486±0.000
Silicon	39.035±0.480	42.540±0.383	51.765±0.345	11.876±0.718
Sodium	63.882±0.000	21.792±0.000	107.568±0.000	13.325±0.000
Zinc	10.688±0.006	15.195±0.006	0.841±0.00	11.363±0.006

Todas as águas subterrâneas contém sais em dissolução em maiores concentrações do que nas águas superficiais, devido a uma maior exposição das mesmas nos extratos geológicos aos materiais solúveis. Os tipos de sais e suas concentrações dependem do meio ambiente, do fluxo e da fonte de água subterrânea. Estes sais originam-se da dissolução dos materiais rochosos pelos quais estas águas percolam.

O conteúdo de sais nas águas subterrâneas deve-se a diversos fatores. A concentração inicial nas águas que escoam no solo para serem infiltradas no aquífero é influenciada por fatores climáticos. Ao infiltrarem as mesmas corroboram para a dissolução dos sais na porção não saturada dos aquíferos (zona de aeração), que serão carregados para a zona saturada dos mesmos. A concentração neste compartimento dependerá dos tipos litológicos dos mesmos e do tempo de escoamento. Evaporação de água próximo a superfície dos mesmos é o fator primordial de enriquecimento em sais das águas subterrâneas<sup>(9)</sup>.

O pH das águas subterrâneas é controlado pelo teor de CO<sub>2</sub> dissolvido nas mesmas e o de carbonatos<sup>(1)</sup>. O pH geral das águas minerais do estado de Santa

Catarina oscila entre 4.6-8.6 e o pH; já o pH das água minerais provenientes de rochas cristalinas varia entre 4.6-7.9<sup>(9)</sup>. O pH das águas estudadas (Tabela 1) corresponde aos valores arbitrados para as águas minerais de Santa catarina e depende do aquífero em que elas permeiam, sendo que a solubilização dos componentes presentes pode alterar o valor. Isto explica a diferença entre as águas subterrâneas do Ponto 1 (águas que permeiam os filões mineralizados) e as do ponto 2 (águas que permeiam as rochas sedimentares), já que as mesmas provém de aquíferos com diferentes composições litológicas (Figure 2)




Figure 2 – Perfil esquemático do Cocal vein e pontos de coleta das águas subterrâneas (P<sub>1</sub> e P<sub>2</sub>).

A água do ponto 3 (lago de decantação) é proveniente da água de lavagem do minério, que provém do Ponto 1. Assim justifica-se a diferença mínima de pH entre estes dois pontos. Os teores de carbonato e bicarbonato estão diretamente relacionados ao pH. Como o pH das águas analisadas provenientes dos pontos da mina encontram-se entre 6.53 e 7.35, isso significa que as águas não apresentam

carbonatos, pois águas com estes ânions tem pH acima de 8.30. Portanto, as águas da Mina 3 apresentam somente bicarbonatos, que se encontram presentes a partir de pH 4.50. Em condições normais a grande maioria das águas que circulam em aquíferos cristalinos apresentam condições de potabilidade sem restrições, com pH entre 6.5 e 7.5, portanto em condições de neutralidade<sup>(10)</sup>. Na Mina 3, os poços com entradas de água acima de 150 m, geralmente apresentam pH entre 6.5 e 7.9, podendo eventualmente alcançarem valores superiores a 8, sempre que os teores de Ca forem inferiores a 10 ppm ou carbonato de cálcio inferiores a 100 ppm. Nos poços onde as entradas de água se deram abaixo ou próximo de 200 m de profundidade, o pH sobe, alcançando em média valores na ordem de 9. Logo, em relação ao pH as água analisadas são neutras.

Em soluções diluídas, como as águas subterrâneas, a condutibilidade elétrica é proporcional à quantidade de minerais dissolvidos nas mesmas, aumentando com o aumento de temperatura<sup>(1)</sup>. A condutividade geral das águas minerais do estado de Santa Catarina varie entre 6.15 e 1.15  $\mu\text{S}\cdot\text{cm}^{-1}$ , o que reflete o grau de mineralização das águas. Os valores de condutividade elétrica nas águas subterrâneas da Mina 3 (Tabela 1) indicam haver intensa atividade iônica no Ponto 1 em relação ao Ponto 2, o que se justifica pelo fato das destas (Ponto 1) estarem em contato direto com o filão de florita, ou seja um aquífero mais rico em fases minerais. Águas altamente mineralizadas podem receber incidência de radioatividade, o que se dá, principalmente, pela presença de radônio<sup>(9)</sup>. Águas que percolam rochas cristalinas tendem a ser mais ricas em minerais radioativos. Assim o alta condutividade elétrica dectada para as águas que circulam pelo Granito Pedras Grandes justifica os altos teores verificados para a condutividade elétrica (Figura 3).

Análises químicas em 80% das águas que circulam aquíferos cristalinos apresentam teor de bicarbonato entre 40 – 80 ppm<sup>(10)</sup>. A presença tanto de carbonato quanto de bicarbonato se reflete diretamente na alcalinidade, que por definição é a capacidade de uma água em neutralizar ácidos. A alcalinidade das águas minerais de Santa Catarina variam de 154.57  $\text{mg}\cdot\text{L}^{-1}$  (129.3 – 189.3) para as

rochas da Formação Serra Geral e  $121.46 \text{ mg.L}^{-1}$  (33.5 – 209.8) para as rochas do embasamento cristalino<sup>(9)</sup>. Nas águas subterrâneas da Mina 3 o teor encontrado para bicarbonato é similar ao encontrado para as rochas cristalinas do estado de Santa Catarina, que podem conter também bicarbonato proveniente das rochas da Formação Serra Geral, que estão presentes também no distrito Fluorítico de Santa Catarina. O bicarbonato é o ânion presente em maior concentração nos pontos 1, 2 e 3. Isto está associado às altas concentrações de Ca e Na, o que pode ser decorrente da solubilização de bicarbonatos. A maior concentração no Ponto 3, significa o incremento do teor durante o processo de lavagem do minério que se acumula no lago de decantação da mina.

O teor de cloreto de uma água subterrânea expressa a quantidade de íons Cl presentes, que podem advir da dissolução de  $\text{CaCl}_2$ , NaCl, KCl e  $\text{KMgCl}_3 \cdot 6\text{H}_2\text{O}$ . Uma água é considerada satisfatória quando o teor de cloreto é menor do que 150 ppm. Portanto, os teores das águas da Mina 3 encontram-se dentro desta faixa.

O íon fluoreto, geralmente, é encontrado nas águas subterrâneas em pequenas concentrações. O mesmo deriva da fluorite ( $\text{CaF}_2$ ), principal halogeneto presente em rochas ígneas, ou então de minerais nos quais o flúor possa estar presente como fluorapatite ( $\text{CaPO}_4)_3\text{F}$ ), topaz ( $\text{Al}_2\text{SiO}_4\text{F}_2$ ), mica and tourmaline. O flúor liberado pelo intemperismo destes minerais ocorre na forma de íon fluoreto, que apresenta alta mobilidade. O mesmo pode formar complexos estáveis com Al, Fe, B e Ca. Assim, no ciclo geoquímico o flúor pode ser removido das águas pela coprecipitação com óxidos secundários de Fe, podendo também ser complexado com Fe e Al na forma de fosfatos. Em águas que percolam litologias cristalinas a presença de flúor é mais significativa em face dos granitoides apresentarem teores mais altos do elemento. Nas águas minerais do estado de Santa Catarina, que percolam o Granito Pedras Grandes, os teores variam de 0,40 – 1,10  $\text{mg.L}^{-1}$ <sup>(9)</sup>. Nas águas subterrâneas da Mina 3 (pontos 1 e 2) (Tabela 2) o teor de fluoreto está acima daqueles das águas minerais do estado de Santa Catarina. Certamente isto se deve à presença de fluorita, pois seus filões cortam as litologias do Distrito Fluorítico de Santa Catarina (Figuras 2 – 3), encaixando-se preferencialmente no Granito Pedras Grandes. A concentração deste ânion no Ponto 1 está acima dos teores indicados por Hausman<sup>(10)</sup>, porém os teores do ponto 2 estão de acordo com

C. Kath, P. C. P. das Neves and L. G. Ionescu

a faixa do 5 ppm<sup>(3)</sup>. O teor desejável de fluoreto nas águas deve estar em torno de 1,00mg.L<sup>-(1-3)</sup>, pois acima deste valor favorece à fluorose, patologia dentária que afeta principalmente crianças, promovendo manchas escuras no esmalte dentário e crescimento protuberante nos dentes incisivos<sup>(11)</sup>. As águas de consumo humano (Ponto 4) apontam 0.42 mg.L<sup>-1</sup>, estando assim dentro dos parâmetros de segurança em relação à fluorose.

Sulfatos em águas subterrâneas são provenientes, principalmente, da presença de gipsita (CaSO<sub>4</sub>.2H<sub>2</sub>O) e anidrita (CaSO<sub>4</sub>), fases minerais que já foram identificadas nas litologias da Formação Serra Geral<sup>(1-12)</sup>. Os teores de sulfato das águas coletadas no Ponto 1, também podem provir da solubilização da pirita, fase mineral encontrada amplamente na Mina 3, associada à fluorita<sup>(6-9)</sup>. O mesmo pode ser aludido à presença do ânion no Ponto 2.

O alumínio encontra-se presente em rochas ígneas, geralmente, na forma de Al<sub>2</sub>O<sub>3</sub>, com teor entre 10-20%<sup>(14)</sup>. Seus íons hidratados são de extrema importância na regulação do pH de águas subterrâneas. Em meios moderadamente ácidos o alumínio disponível, na forma iônica é convertido em hidróxido de alumínio com estruturas que podem variar de Al(OH)<sup>2+</sup> a [Al<sub>10</sub>(OH)<sub>22</sub>]<sup>8+</sup>, processo que ocorre concomitante com outros cátions como o ferro e o manganês. Deste modo pode ocorrer alterações do pH do meio. A intensidade do processo está diretamente relacionada com as quantidades do elemento presentes nas formas disponíveis, além da intensidade da interação entre o corpo aquoso e o leito pelo qual flui. As concentrações de alumínio em águas subterrâneas podem variar entre 100mg.L<sup>-1</sup> a 2000mg.L<sup>-1(13)</sup>. O alto teor de alumínio nas águas de rejeito (P<sub>3</sub>) (tabelas 3-4) pode ser atribuído, em parte, as interações entre a água subterrânea e os sedimentos. O aumento da concentração de alumínio no ponto 3 em relação a águas do ponto 2, contraria a teoria do aumento da sua concentração em função do valor do pH, já que estas águas apresentam praticamente os mesmos pH. Isto pode significar que algum outro fator, ou elemento químico, esteja sendo mais determinante. As concentrações do elemento presentes nas águas da Mina Fluoral em termos de concentração total

(Tabela 4), podem indicar que parte do alumínio presente neste meio não esteja disponível. O teor de alumínio presente no ponto 2 encontra-se dentro da faixa indicada por Allen (1974)<sup>(13)</sup>. Já, as águas do ponto 1, encontram-se acima desta faixa. Isto pode estar relacionado à composição da rocha que forma o aquífero, podendo a mesma conter minerais com o elemento em sua constituição, ou, ainda, estes minerais serem muito pouco solúveis.

O cálcio é o quinto elemento químico mais abundante na crosta terrestre, com 3.63%<sup>(14)</sup>. As principais fontes do elemento são os plagioclásios cálcicos, calcite, dolomite, fluorite, apatites and anfíboles. Na forma de carbonato de cálcio é pouco solúvel em água. Ocorre, portanto, nas águas na forma de bicarbonato e sua solubilidade se dá na quantidade de gás carbônico dissolvido. Nas águas subterrâneas as variações de temperatura e pressão modificam o CO<sub>2</sub> dissolvido, o que reflete sobre o conteúdo em cálcio. Estas variações, ora levam à solubilização do carbonato de cálcio, ora levam a sua precipitação. Segundo Allen (1974)<sup>(13)</sup>, em águas naturais o cálcio encontra-se na faixa de 1 a 100mg.L<sup>-1</sup>. O teor em cálcio nas águas do ponto 1 (Tabelas 3 e 4) pode estar relacionado com a presença de fluorite, que é o principal mineral constituinte do filão por onde estas águas percolam. A diferença entre a concentração de cálcio total e disponível indica que grande parte dele não está solubilizado, podendo estar precipitado ou associado a outros elementos. A atividade e disponibilização do cálcio são favorecidos numa faixa de pH entre 6,5-8,5 (Brady, 1979)<sup>(13)</sup>. Assim, podee ser justificada a diferença na concentração de cálcio entre os pontos 1 e 2, que também pode estar associada à composição mineral do aquífero correspondente às águas do ponto 2, que percolam rochas sedimentares.

O ferro, quarto elemento em abundância na crosta terrestre, com 5.00%<sup>(14)</sup>, está presente em todas as águas potabilizáveis. O ferro comumente se encontra presente nas águas na forma de Fe<sup>2+</sup> e Fe<sup>3+</sup>. O íon ferroso é instável em presença do ar, mudando para o estado férrico quando a água que o contém é exposta, sendo que pode dissolver-se em quantidade de cerca de 50ppm, em águas neutras e em maiores proporções se as águas forem levemente ácidas. Já o íon férrico tende à insolubilidade em águas alcalinas ou fracamente ácidas. O alto teor de ferro nas águas do ponto 3 pode estar relacionado ao contato da mesma com as tubulações

C. Kath, P. C. P. das Neves and L. G. Ionescu

da Mina, já que a mesma é utilizada para a lavagem do minério. A diferença entre a quantidade disponível e a total, pode significar que parte do ferro está na forma insolúvel, o que pode ter sido causado, também, pela ação de microorganismos presentes no lago de decantação ou pelo contato com o ar, podendo estar na forma de óxidos insolúveis. Segundo Hausman<sup>(10)</sup>, os teores mais elevados de Fe ocorrem quando as águas dos granitóides são contaminadas pelas águas que circulam pelas formações sedimentares sobrepostas. Isto pode justificar a diferença entre os teores de ferro do ponto 1 e do ponto 2 (Tabelas 3 e 4). O ferro nas águas do ponto 1 pode ter a sua fonte principal nos minerais goethite ( $\alpha\text{-Fe}^{3+}\text{O(OH)}$ ), hematite ( $\alpha\text{-Fe}_2\text{O}_3$ ) e pyrite ( $\text{FeS}_2$ ), presentes na composição rochosa do aquífero<sup>(6)</sup>. Concentrações entre 1-5  $\text{mg.L}^{-1}$  de ferro em águas subterrâneas são comuns<sup>(1)</sup>. Após a aeração, o teor pode cair a 0.1  $\text{mg.L}^{-1}$ . O teor disponível deste elemento nos pontos 1 e 2 (Tabela 3) encontra-se abaixo da faixa indicada, podendo ter sido reduzido pela aeração provocada quando do bombeamento da água do aquífero até a superfície.

O lítio na crosta terrestre é um elemento traço (20 ppm/Kg)<sup>(14)</sup>, o que justifica sua baixa concentração nas águas subterrâneas. Nas amostras dos pontos 1, 2 e 3, os teores de lítio demonstram comportamento diretamente proporcional às variações de pH. A pequena disparidade entre os valores para as concentrações do elemento nas formas solúvel e total, nos pontos 1 e 2 (Tabelas 3 e 4), supõe que praticamente todo o lítio está na forma solúvel. No ponto 3 a concentração praticamente dobrou da forma solúvel para a total, o que significa que boa parte do mesmo não está disponível para o meio, podendo em consequência ter ficado retida nos sedimentos.

O magnésio tem uma participação de 2.09% na crosta terrestre<sup>(14)</sup>. Seu comportamento geoquímico é semelhante ao do cálcio, com a diferença de que forma sais mais solúveis. Encontra-se presente em minerais como a biotite, anfíboles, olivines e piroxenes, mais estáveis ao intemperismo químico do que aqueles fornecedores de cálcio, por isso a presença de magnésio é menor do que a de cálcio nas águas subterrâneas. A atividade química e a disponibilidade do

magnésio está diretamente ligada ao pH do meio, decrescendo em função da diminuição do mesmo. Assim, em meios mais básicos ou neutros sua disponibilidade tende a ser mais alta. Ávila et al. <sup>(9)</sup> atribuíram valor de 0.96 mgL<sup>-1</sup> para aquíferos da Serra Geral Formation (basaltos), 2.28 mgL<sup>-1</sup> para aquíferos do Pedras Grandes Granite 2 e 28.33 mgL<sup>-1</sup> para as águas da Rio do Sul Formation. Os valores encontrados para o ponto 1, estão de acordo com os valores acima, podendo parte do magnésio encontrado ser proveniente da Rio do Sul Formation, presente junto ao Pedras Grande Granite no Fluoritic District of Santa Catarina. O maior teor de Mg no ponto 2 (Figura 3) pode ser função de sua maior proximidade com a superfície, onde há maior dissolução. A diferença entre a quantidade disponível e a total (Tabelas 3 – 4) pode ser atribuída à facilidade deste elemento formar sais mais solúveis, levando à ilação de que praticamente todo o magnésio esteja solubilizado.

O manganês ocorre na crosta terrestre com teor de 0.09% <sup>(14)</sup>. Embora, quimicamente, seu comportamento assemelha-se ao do ferro em ocorrência nas águas naturais, seus teores são sensivelmente menores devido a sua ocorrência crustal ser menor do que a daquele elemento. Nos pontos 1, 2 e 3 (Tabelas 2 e 4) do local estudado, o Mn é o elemento de menor concentração. Segundo Hausman <sup>(10)</sup>, em águas que percolam o cristalino os teores em Mn são da ordem de 0.05 ppm, ficando assim os valores obtidos nos pontos 1 e 2 deste estudo inseridos na faixa citada.

O silício é o segundo elemento mais abundante da crosta terrestre (27.72%) <sup>(14)</sup>. Combinado com o oxigênio, forma o mineral mais difundido na natureza (quartz). Segundo a CETESB <sup>(1)</sup>, a água pode dissolver somente diminutas quantidades de sílica; não obstante as águas subterrâneas podem conter até 100 ppm do óxido., sendo que valores entre 0,5-8 mg.L<sup>-1</sup> são encontrados para o silício nas águas subterrâneas <sup>(13)</sup>. Segundo Mason <sup>(14)</sup>, em pH menor que 4 a sílica dissolve lentamente, entretanto entre 5-9 sua solubilidade tende a aumentar consideravelmente. Ávila et al. <sup>(9)</sup> afirmam que o fator determinante para a variação do teor em sílica nas águas subterrâneas depende da litologia, clima e pH. Nas rochas cristalinas de Santa Catarina (Pedras Grandes Granite) está relacionada com a decomposição química dos minerais silicatados (feldspates,

C. Kath, P. C. P. das Neves and L. G. Ionescu

plagioclases e quartz). Nas águas dos pontos 1, 2 e 3 o Si é o segundo cátion com maior concentração nas águas. O fato de que grande parte dele não se encontra solubilizado deve-se a sua grande resistência a sua desintegração geoquímica (elemento altamente estável). A alta concentração registrada nos meios estudados (Tabela 3), pode ter sido facilitada pelo pH das águas, que se encontram dentro da faixa de solubilidade da sílica. Os teores aqui encontrados estão em consonância com os de Ávila et al.<sup>(9)</sup> para os do Pedras Grandes Granite.

O sódio e o potássio ocorrem com distribuição geoquímica muito próximas na crosta terrestre, com 2.83 e 2.59 %, respectivamente<sup>(14)</sup>, muito embora sua participação nas águas subterrâneas se dá em quantidades sensivelmente diferentes. O potássio ocorre na estrutura de muitos minerais (micas, feldspates, sylvite), entretanto pode ficar retido nas argilas o que o torna menos disponível para o meio. Sua faixa de concentração em águas subterrâneas se dá na ordem de 0.5 a 10 mg.L<sup>-1</sup>, logo as águas dos pontos 1 e 2 estão de acordo com esta faixa. Nas águas de litologias cristalinas o K ocorre com melhor representatividade, pois as mesmas são muito ricas em feldspates<sup>(9)</sup>. Segundo Todd<sup>(15)</sup>, as águas subterrâneas que percolam rochas ígneas dissolvem pequenas quantidades de substâncias minerais, devido a relativa insolubilidade da composição destas rochas. Isto pode justificar o alto teor de K na forma total das águas do ponto 1 em relação à forma solúvel deste mesmo ponto, e também, à pequena diferença entre o teor disponível e o teor total na amostragem do ponto 2, já que o aquífero por onde circulam estas águas é constituído por rochas sedimentares que são mais solúveis do que as ígneas. O sódio é um elemento químico quase sempre presente nas águas subterrâneas e, deriva de uma série de minerais (feldspates, anfíboles e halite). Por ser muito solúvel em ambientes minerotróficos há uma grande tendência à perda do mesmo por solubilização. Sua atividade não sofre mudanças em função de alterações de pH. Sais de Na formados em processos intempéricos são geralmente muito solúveis. Os valores médios para o Na em águas naturais varia entre 2 – 100 mg.L<sup>-1</sup><sup>(13)</sup>. Nos pontos 1, 2 e 3 das águas da Mina Fluoral observou-se que o Na é o cátion com a maior concentração. Isto se justifica pela alta solubilidade do

elemento. O alto teor do íon em concentração total (pontos 2 e 3) – (Tabela 4), significa que grande parte do mesmo não está solubilizado. Segundo Ávila et al.<sup>(9)</sup>, as águas da Serra Geral Formation são as que apresentam os maiores teores do elemento, devido ao fato destas rochas serem muito ricas em feldspates e pyroxenes, minerais ricos em Na.

O zinco participa da composição da crosta terrestre como elemento-traço (70 ppm)<sup>(14)</sup>. Os principais minerais portadores do elemento são sphalerite (ZnS), zincite (ZnO) e hemimorphite ( $Zn_4Si_2O_7(OH)_2 \cdot H_2O$ ), não registrados para a Mina Fluoral. O baixo teor de Zn nas águas do lago de decantação (ponto 3), tanto na forma solúvel, quanto na total (Tabelas 3 - 4), pode ser atribuído ao fato de que praticamente todo o zinco presente nas águas do ponto 1 está na forma residual insolúvel, muitas vezes como colóide, tendo se depositado no fundo do lago. Também. Parte do mesmo pode ter sido complexado pelos microorganismos, plantas e matéria orgânica presentes neste ambiente. A alta concentração de Zn na forma total para as águas dos pontos 1 e 2, pode ser explicada por carreamento advindo das tubulações da Mina, pois os minerais de Zn não ocorrem nas litologias locais.

#### **CLASSIFICAÇÃO HIDROGEOQUÍMICA DAS ÁGUAS SUBTERRÂNEAS DA MINA FLUORAL**

A classificação das águas subterrâneas foi efetivada através do diagrama trilinear de Piper<sup>(10)</sup> para as concentrações de  $Ca^{2+}$ ,  $Mg^{2+}$ ,  $K^+$ ,  $HCO_3^{4-}$ ,  $CO_3^{2-}$ ,  $Cl^-$  e  $SO_4^{2-}$  em  $Meq.L^{-1}$  e para os valores de condutividade elétrica nas quatro amostras analisadas. As águas do ponto 1 foram classificadas como cloretadas sódicas; as do ponto 2 como bicarbonatadas calco-sódicas; as do ponto 3 como bicarbonatadas sódicas.

#### **CONCLUSÕES**

No presente estudo foram estudados apenas os elementos de maior importância e presentes em quantidades mais significativas na constituição

C. Kath, P. C. P. das Neves and L. G. Ionescu

mineral dos aquíferos da Mina Fluoral (Mina 3), Morro da Fumaça, Santa Catarina, Brazil.

- O pH das águas ficou dentro de uma faixa considerada neutra;
- os elementos químicos presentes nas águas subterrâneas, em geral, apresentaram atividades diretamente relacionadas às composições minerais dos aquíferos por onde estas águas circulam;
- o aumento de profundidade de circulação das águas subterrâneas fez aumentar os teores de mineralizações das mesmas, pois à medida que aumenta a profundidade, aumenta o volume de rochas mineralizadas;
- a classificação hidrogeoquímica das águas estudadas demonstrou haverem cloradas sódicas, águas bicarbonatadas sódicas e águas bicarbonatadas calcosódicas na Mina Fluoral;
- a maior concentração dos elementos químicos analisados deu-se no ponto 3, o que se justifica por serem águas de lavagem do minério, ocorrendo um incremento pela dissolução do mesmo e, também, porque no lago não há movimentação da água, o que facilita a sua concentração, além da interação com os sedimentos e os microorganismos existentes no mesmo;
- a maior concentração de cátions na forma total, nas águas do ponto 1 em relação às do ponto 2, é explicada pelo fato de que as águas subterrâneas que passam por rochas ígneas dissolvem pequenas quantidades de substâncias minerais, ao contrário das rochas sedimentares, que são mais solúveis;
- pela comparação das águas do ponto 2 e do ponto 4, juntamente com padrões estabelecidos pelo CONAMA, conclui-se que as águas do ponto 2 apresentam menores condições de potabilidade;
- os valores encontrados nas águas do ponto 1 estão de acordo com os obtidos para as águas minerais do Estado de Santa Catarina, ambas circundantes do Pedras Grandes Granite.

Ao serem analisadas as características hidrogeoquímicas das águas que percola o embasamento cristalino de uma região, deve-se sempre levar em consideração as anomalias relacionadas à contribuição de certos elementos, pois

fatores de ordem tectônica podem induzir, devido a mineralizações, variações na qualidade das águas, elevando em consequência os teores de alguns elementos químicos.

#### REFERENCES

- (<sup>1</sup>) CETESB – Companhia de Tecnologia de Saneamento Ambiental do Estado de São Paulo. *Água subterrânea e poços tubulares*. 2<sup>a</sup> ed., São Paulo. In: Johnson Division, Minnesota, 1974, 292p.
- (<sup>2</sup>) GLEICK, P. H. *World's water 2004-2005: the biennial report on freshwater resources*. Island Press, 2005.
- (<sup>3</sup>) CONAMA – Qualidade da água. Resolução CONAMA n<sup>o</sup> 396 de 03.04.2008, 2008 - DOU n<sup>o</sup> 66, 07.04.2008, seção I, pp. 64-68, 2008.
- (<sup>4</sup>) SALLET, R. *Étude pétrologique et métallogénique d'un secteur du district à fluorine de Santa Catarina, Brésil. Les granitoides précambriens monzonitiques source probable de la fluorine filonienne post-jurassique*. Paris, 233p. Thèse de Doctorat, Université de Paris VI., 1988.
- (<sup>5</sup>) TEIXEIRA, C. A. S. Relatório de datações geocronológicas do projeto básico da região leste de Santa Catarina e do projeto específico da fluorita. *Boletim do Departamento Nacional da Produção Mineral*, 1<sup>o</sup> Distrito, Porto Alegre, 39p., 1969.
- (<sup>6</sup>) BASTOS NETO, A. C.; JELINEK, R. Uma ocorrência singular de barita do sudoeste catarinense, Brasil. Porto Alegre: *Pesquisas*, v. 24, n<sup>os</sup>. 1-2, 31-38, 1997.
- (<sup>7</sup>) BACK, M. E. *Fleischer's glossary of mineral species*. Tucson: The Mineralogical Record, 2014, 434p.
- (<sup>8</sup>) STANDARD METHODS FOR THE EXAMINATION OF WATERS AND WASTEWATERS. American Public Health, 864p., 1981.
- (<sup>9</sup>) ÁVILA, E. L.; AWDZIEJ, J.; BASTOS, J. P. C.; PARAHYBA, R. E. R.; BICCA, V. H. F. Águas minerais do estado de Santa Catarina. *Acta Geologica Leopoldensia*, São Leopoldo, XIII(3), 197-212, 1990.
- (<sup>10</sup>) HAUSMAN, A. Províncias hidrogeológicas do Estado do Rio Grande do Sul. *Acta Geologica Leopoldensia*, São Leopoldo, série mapas, mapa n. 2, 127p., 1995.

C. Kath, P. C. P. das Neves and L. G. Ionescu

- (11) SATAKE, A.; ZAIA, W. L. S.; COUTINHO, M. Remoção de manchas fluoróticas através da técnica de microabrasão: relato de caso clínico. *Revista Dental Press Estética*. Maringá, vol. 7, n. 1, p. 118-125, 2010.
- (12) NEVES, P. C. P. das; ATENCIO, D. *Enciclopédia dos minerais do Brasil*. ULBRA: Canoas, 431p., 2014.
- (13) PROCHNOW, E. A. Caracterização dos parâmetros geoquímicos e bioindicadores de um solo orgânico na Fazenda São Maximiano, Guaíba, Rio Grande do Sul, Brasil. Canoas, 170p. Monografia de Bacharelado em Química, ULBRA, 123f., 2001.
- (14) KLEIN, C; DUTROW, B. *Manual de Ciência dos Minerais*. Porto Alegre: Bookmann, 2012, 710p.
- (15) TODD, D. K.. *Hidrologia das águas subterrâneas*. São Paulo: Edgard Blücher, 1959, 319p.

**Volatile Components of *Oncoba spinosa* leaf and *Morus mesozygia* leaf and stem bark**

S. A. Aboaba\* and O. Adeeko<sup>1</sup>

*Department of Chemistry, University of Ibadan, Ibadan, Nigeria*

G. Flamini<sup>2</sup>

*Dipartimento di Farmacia, via Bonanno 33, 56126, Pisa Italy*

\* Corresponding author: [saboaba@gmail.com](mailto:saboaba@gmail.com)

1 [kemiadeeko@yahoo.com](mailto:kemiadeeko@yahoo.com)

2 [guido.flamini@farm.unipi.it](mailto:guido.flamini@farm.unipi.it)

**ABSTRACT**

The essential oils of *Oncoba spinosa* leaf and *Morus mesozygia* leaf and stem bark were extracted using hydro distillation and analyzed by means of Gas chromatography (GC) and GC coupled with mass spectrometry (GC-MS). The yields of the essential oils were; 0.50 %, 0.165 % and 0.456 % respectively for *Oncoba spinosa* leaf, *Morus mesozygia* leaf and stem oil. A total of twenty eight, thirty four and twenty compounds representing 92.0%, 92.0% and 96.9% of the total oil contents were identified, respectively from the leaf of *O. spinosa*, leaf and stem oil of *M. mesozygia*. Leaf oil of *O. spinosa* contained linalool (22.1 %),  $\beta$  – caryophyllene (18.7 %), caryophyllene oxide (10.6 %) and pentadecanal (5.6 %) as the main constituents. *M. mesozygia* leaf oil was dominated with  $\beta$  – elemene (11.7 %), (*E*) – $\beta$ - ionone (12.4 %),  $\alpha$ - selinene (5.1 %), germacrene A (6.0 %),  $\delta$  – cadinene (4.7 %) and spathulenol (7.4 %) while *M. mesozygia* stem oil had 2 –dodecanone (77%) and hexahydrofarnesylacetone (13 %) as its main constituents.

**Key words:** *Oncoba spinosa*, *Morus mesozygia*, linalool, (*E*) – $\beta$ - ionone, 2 –dodecanone.

## INTRODUCTION

Plants essential oils and their components have been known to exhibit biological activities. Screening of plants extracts or their essential oil for these properties has become of great importance because of their growing interest in food and pharmaceutical industries (1). *Oncoba spinosa*, commonly known as the snuff-box tree is a plant species in the genus *Oncoba* and family *Flucourtiaceae*. It is a small to medium-sized deciduous tree of about 5 m high and has simple leaves. The tree is widely distributed along the eastern side of Africa as far as South Africa, mainly in dry woodland or open savanna in a wide range of sites from river valleys to rocky hills. In African medicine, the roots are used in the treatment of dysentery and bladder complaints and fruits in the treatment of leprosy. The anti-ulcerogenic potential (2), antihyperglycemic and antineoplastic activities (3), and anti-convulsant property (4) have been studied. The fruit was found to be effective in the treatment of urinary tract diseases of cows and camels (5). In addition, the antipasmodial (6) and antihelminthic properties (7) have been discussed in literature.

*Morus mesozygia* (Moraceae), commonly called black mulberry or African mulberry, is a small to medium sized forest tree. *Morus* comprises of ten to fifteen species with only one specie (*Morus mesozygia*) native to tropical Africa. It has a wide distribution in tropical Africa, from Senegal eastward to Ethiopia and southward to Zambia, Angola, Mozambique and South Africa (8). In African traditional medicine, parts of *M. mesozygia* are used in decoctions, baths, massages and enemas against rheumatism, lumbago, intercostal pain, neuralgia, colic, stiffness,

debility, diarrhea and dysentery. It is evident from literature that extract of the plant is known to have antidepressant property (9). Arylbenzofurans and flavonoids were isolated from *M. mesozygia* (10) and the anti-plasmodial and cytotoxic properties of arylbenzofuran have been studied (11). The leaf essential oil of another species *Morus rotundiloba* was characterized and dominated by benzyl alcohol (10.48%), dihydroactinolide (10.11%) and palmitic acid (7.01%); the oil also displayed good antibacterial and cytotoxic activities (12).

Therein, we describe the chemical composition of the essential oil from the leaves of *O. spinosa* and leaves and stem bark of *M. mesozygia* for the first time.

## EXPERIMENTAL

### Plant materials

Plant samples were collected from the Botanical garden, University of Ibadan, Ibadan, Nigeria in March 2014 by Mr. Owolabi. The plants were authenticated at the Herbarium of the Forestry Research Institute of Nigeria (FRIN) Oyo State, Nigeria where voucher specimens FHI 110037 and FHI 110038 for *Oncoba spinosa* and *Morus mesozygia* respectively were deposited. Plant samples were air-dried prior to extraction.

### Isolation of essential oils

250 g of each of the dried plant samples were shredded and their oils were obtained by separate distillation for 3 hours in a Clevenger type apparatus. The essential oil fractions obtained were kept in a refrigerator at 4°C prior to analysis.

**Analysis of the Oils**

Gas chromatography (GC) analysis was performed on a HP-5890 Gas chromatograph equipped with a Flame ionization detector and fitted with HP-5 capillary column (30 m x 0.25 mm, 0.25  $\mu$ m film thickness). The GC oven temperature was programmed at 60°C (held for 10 min), heated to 220°C at 5°C/min. The injector and detector temperatures were maintained at 250°C. Helium was used as carrier gas at a flow rate of 2 mL/min with a split ratio of 1:30.

Gas chromatography / Mass spectrometry (GC-MS) analysis were carried out on a Varian CP-3800 gas chromatograph interfaced to a Varian Saturn 2000 ion trap Mass Detector operated at 70 eV. The injector and transfer line temperatures were 220°C and 240°C, respectively. The GC oven temperature was programmed from 60 to 240°C at 3°C/min. Helium was used as a carrier gas at a flow rate of 1 mL/min, injection of 0.2  $\mu$ L aliquots as 10% Hexane solution, split ratio of 1:30.

Identification of the constituents on both columns was based on comparison of the retention times with those of authentic samples, comparing their retention indices relative to the series of n-hydrocarbons, and by comparison of their mass spectra with published spectra and those of reference compounds from NIST 98 and ADAMS (13, 14, 15, 16, 17). The relative concentration of each constituent was calculated by integration of GC peak areas.

**RESULTS AND DISCUSSION**

A total of 28 compounds amounting to 92% were identified in *O. spinosa* leaf essential oil. Among these are, 29.2% oxygenated monoterpenes, 22.3% sesquiterpenes and 9.2% oxygenated sesquiterpenes. It also contained non-terpene derivatives (14.7%) and apocarotenes

(16.6%). However, the major constituents identified are; linalool (22.1%),  $\beta$ - caryophyllene (18.7%), caryophyllene oxide (10.6%), pentadecanal (5.6%), methyl salicylate (4.2%) and (*E*) -  $\beta$ -ionone (3.8%). Linalool and  $\beta$ - caryophyllene which dominated the oil are noted for many commercial and medicinal applications. Linalool, a naturally occurring terpene alcohol has been utilized in perfumery (18) while  $\beta$ - caryophyllene is a dietary cannabinoid (19) which has been found to have clearly demonstrated its ability in the regulation of emotional behaviour, thereby suggesting that its receptor could be relevant therapeutic target for the treatment of anxiety and depressive disorders (20).

A total of 34 constituents were identified in the leaf of *M. mesozygia*, representing 92.0% of the total fraction. Monoterpenes (0.5%), sesquiterpenes (48.5%), oxygenated sesquiterpenes (22.1%), apocarotenes (18.4%) and non-terpene derivatives were identified in the essential oil. The main constituents are; (*E*)- $\beta$ -ionone (12.4%),  $\beta$ -elemene (11.7%), Spathulenol (7.4%),  $\alpha$ -selinene (5.1%),  $\delta$ -cadinene (4.7%), humulene epoxide II (3.8%) (*E*) -  $\alpha$ -ionone (3.4%), trans- $\alpha$ -bergamotene (3.5%). (*E*)- $\beta$ -ionone is a flavouring agent and exhibits anti-proliferative function (21) while  $\beta$ -elemene has exhibited tremendous anti-tumor properties (22, 23).

The essential oil of the stem bark of *M. mesozygia* afforded identification of 20 constituents, corresponding to 96.9% of the oil constituents. Monoterpenes (0.1%), oxygenated monoterpenes (0.5%), sesquiterpene (2.5%), oxygenated sesquiterpenes (1.4%), apocarotenes (14.0%) and non-terpene derivative (78.4%) were identified in the essential oil. The main constituents in the oil are; 2-dodecanone (77.0%), hexahydrofarnesyl acetone (13.0%), ar-curcumene (1.0%). Moreover, it was observed that both *M. mesozygia* leaf and stem bark essential oils had *E*-  $\beta$ - ionone (12.4% and 0.4%) respectively as common constituents and also the most abundant in the leaf oil of *M. mesozygia*. Other common constituents are spathulenol

**S. A. Aboaba, O. Adeeko and G. Flamini**

(7.4% and 0.3%), caryophyllene oxide (2.9% and 0.5%) alloaromadendrene (1.5% and 0.3%), *E*-geranylacetone (2.1% and 0.6%),  $\alpha$ -humulene (1.1% and 0.2%),  $\beta$ -caryophyllene (1.2% and 0.6%), limonene (0.5% and 0.1%) and 2-pentylfuran (0.5% and 0.3%) respectively. The dominance of the stem oil of *M. mesozygia* by 2-dodecanone and hexahydrofarnesyl acetone has indicated that the bark oil may exhibit excellent insect repellent properties. In a study by Ndungu et al (24), it was revealed that 2-dodecanone was one of the constituents in the oil of *Cleome monophylla* that exhibited high repellent activity against tick and maize weevil. In another related work, it was deduced that 2-dodecanone exhibited good protection efficacy against *An. gambiae* (100% protection at 10% concentration) and also concluded that methyl ketone analogies are potential source of insect repellent including mosquitoes (25). Hexahydrofarnesylacetone is also a widespread ketone found in plant and insects which several insect uses as part of their pheromone bouquet (26).

The abundance of both compounds 2-dodecanone and hexahydrofarnesylacetone totaling 90% of the stem oil of *M. mesozygia* gives a strong lead that it possesses good insect repellent properties and has therefore opened an avenue to research into the repellent ability of the essential oil to different species of insects. On the contrast, these compounds were not found in the leaf oil yet confirming a known fact that plant tissues differ in their chemical constituents. The constituents of these oils were at variance to the compounds characterized in the oil of another species *Morus rotundiloba*, fifty-one compounds were detected in the leaf essential oil with benzyl alcohol (10.48%), dihydroactinolide (10.11%) and palmitic acid (7.01%) being the most abundant (12). Out of the compounds characterized in *M. rotundiloba*, only four were found in

either of the two oils of *M. mesozygia*, these are E- $\alpha$ -ionone, hexahydrofarnesylacetone, geranylacetone and  $\beta$ -cyclocitral.

### CONCLUSION

The different therapeutics and commercial importance of the characterized compounds in these essential oils justifies the use of different parts for the treatment of different ailments. This represents the first attempt to our knowledge at characterizing the essential oil constituents of the leaves of *O. spinosa* and leaves and stem bark of *M. mesozygia*.

**Table 1: Essential Oil Constituents of the Leaves of *Oncoba spinosa* and Leaves and stem bark of *Morus mesozygia*.**

S/N	Constituents	L.R.I <sup>a</sup>	L.R.I <sup>b</sup>	<i>O. spinosa</i> Leaves	<i>M. mesozygia</i> Leaves	<i>M. mesozygia</i> Stem bark
1	(E) -2-hexenal	865	846	0.8	-	-
2	p-xylene	867	883 <sup>a</sup>	-	0.5	-
3	6-methyl-5-heptene-2-one	987	992	-	-	0.4
4	2-pentyl furan	993	984	0.7	0.5	0.3
5	(E) -2-pentenyl furan	1004	1000 <sup>b</sup>	0.7	-	-
6	Limonene	1032	1024	-	0.5	0.1
7	1,8-cineole	1034	1026	-	-	0.3
8	Linalool	1101	1095	<b>22.1</b>	-	-
9	Nonanal	1103	1100	-	0.5	-

S. A. Aboaba, O. Adeeko and G. Flamini

10	$\alpha$ -terpeneol	1191	1186	3.1	-	-
11	Methyl salicylate	1192	1190	4.2	0.5	-
12	$\beta$ -cyclocitral	1222	1217	0.9	0.5	-
13	Nerol	1230	1227	1.2	-	-
14	Methyl carvacrol	1246	1298	-	-	0.2
15	Geraniol	1256	1249	2.8	-	-
16	$\alpha$ -copaene	1377	1374	0.7	2.9	-
17	$\beta$ -elemene	1392	1389	0.9	<b>11.7</b>	-
18	2-dodecanone	1395	1394	-	-	<b>77.0`</b>
19	<i>cis</i> - $\alpha$ -bergamotene	1416	1411	-	0.5	-
20	$\beta$ -caryophyllene	1419	1417	<b>18.7</b>	1.2	0.6
21	$\alpha$ -santalene	1420	1416	-	1.3	-
22	( <i>E</i> )- $\alpha$ -ionone	1428	1428	1.1	3.4	-
23	Trans- $\alpha$ -bergamotene	1438	1432	-	3.5	-
24	Precocene1	1448	1440 <sup>d</sup>	-	-	0.3
25	epi- $\beta$ -santalene	1450	1445	-	0.5	-
26	$\alpha$ -humulene	1455	1452	1.1	1.1	0.2
27	( <i>E</i> )-geranylacetone	1455	1453	1.5	2.1	0.6
28	Alloaromadendrene	1462	1461	-	1.5	0.3
29	Cabreuva oxide B	1463	1462	0.8	-	-
30	$\gamma$ -muurolene	1478	1478	-	1.8	-

# SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY

SOUTH. BRAZ. J. CHEM., Vol. 23, No. 23, 2015

*Vol. Compo. of O. spinosa leaf and M. mesozygia leaf and stem bark*

31

31	Germacrene D	1482	1484	-	1.3	-
32	ar-curcumene	1483	1479	-	-	1.0
33	(E)- $\beta$ -ionone	1487	1487	3.8	<b>12.4</b>	0.4
34	10,11-epoxycalamenene	1492	1476 <sup>e</sup>	-	-	0.4
35	$\alpha$ -selinene	1496	1498	-	5.1	-
36	Germacrene A	1504	1508	-	6.0	-
37	$\beta$ -bisabolene	1509	1505	-	2.0	0.3
38	$\beta$ -curcumene	1513	1514	0.9	2.3	
39	$\delta$ -cadinene	1524	1513	-	<b>4.7</b>	-
40	(E)- $\gamma$ -bisabolene	1534	1529	-	0.5	-
41	Dihydroactinolide	1536	1531 <sup>f</sup>	1.9	-	-
42	$\alpha$ -calacorene	1543	1544	-	0.6	-
43	(E)-nerolidol	1565	1561	1.8	-	-
44	Caryophyllene alcohol	1569	1568	-	-	0.6
45	(Z)-3-hexenyl benzoate	1570	1564 <sup>g</sup>	0.5	-	-
46	Spathulenol	1577	1577	0.7	<b>7.4</b>	0.3
47	Caryophyllene oxide	1582	1582	<b>10.6</b>	2.9	0.5
48	Viridiflorol	1591	1592	1.1	-	-
49	Humlene epoxide II	1607	1608	0.8	3.8	-
50	Tetradecanal	1612	1611	1.2	-	-

S. A. Aboaba, O. Adeeko and G. Flamini

51	Humulene-1,6-diene-3-ol	1618	1619 <sup>h</sup>	-	1.9	-
52	Cedrenol	1630	1642 <sup>l</sup>	-	1.7	
53	Caryophylla-4(14),8(15)-dien-5-ol	1637	1623 <sup>j</sup>	1.1	-	-
54	Selin-11-en-4- $\alpha$ -ol	1655	1658	-	1.9	-
55	$\alpha$ -bisabolol	1684	1685	-	0.7	-
56	Pentadecanal	1716	1711 <sup>k</sup>	<b>5.6</b>	-	-
57	Hexahydrofarnesyl acetone	1845	1843 <sup>l</sup>	-	-	<b>13.0</b>
58	Benzyl salicylate	1864	1864	-	-	0.4
1	Monoterpene hydrocarbons			0.0	0.5	0.1
2	Oxygenated monoterpenes			29.2	0.0	0.5
3	Sesquiterpenes hydrocarbons			22.3	48.5	2.5
4	Oxygenated sesquiterpenes			9.2	22.1	1.4
5	Apocarotenes			16.6	18.4	14.0
6	Non-terpene derivatives			14.7	2.5	78.4
	Total Identified			<b>92.0</b>	<b>92.0</b>	<b>96.9</b>

“Notes”

L.R.I<sup>a</sup> = Linear Retention Index From This Work

L.R.I<sup>b</sup> = Linear Retention Index From Literature (Adams)

<sup>h-l</sup> = References 27-35.

## REFERENCES

1. P.H Amvam Zollo, L. Biyiti, F. Tchoumboungang, C. Menut, G. Lamaty and P. Bouchet, *Flav. Fragr. J.*, 13: 107-114 (1998).
2. M. Olaleye and O. Titilope, *Global J. Pure Appl. Sci.*, 12(3): 369-371 (2006).
3. O.S Balogun, I.A. Oladosu, A. Akinnusi and L. Zhiqiang, *Elixir Appl. Chem.*, 59: 15637-15641 (2013).
4. J. Risa, A. Risa, A. Adsersen, B. Gauguin, G. I. Stafford, J. Van Staden, A.K. Jäger, *J. Ethnopharmacol.*, 93: 177-182 (2004).
5. A.N. Mohammed, H. Sher and L. Wijaya, *J. Med. Plt. Res.*, 4(21): 2298-2304 (2010).
6. I.T. Gbadamosi and A.A. Oloyede, *Afr. J. Pharm. & Pharmacol.*, 8(23): 638-643 (2014).
7. R.A. Mothana, S. Kriegisch, M. Harms, K. Wende and U. Lindequist, *Pharm. Biol.*, 49(2): 200-210 (2011).
8. H.M. Burkill, *The Useful Plants of West Tropical Africa, 2<sup>nd</sup> Ed. Vol. 4. Families M-R. Royal Botanical Gardens, Kew, Richmond, U.K.*, pp 969 (1997).
9. F. Adediwura and O. Bola, *Pharmacol. & Pharm.*, 4(1): 100-103 (2013).
10. G.D.W.F. Kapache, D. Amadou, P. Waffo-Teguo, J.H. Donfack, C.D. Fozing, D. Harakat, A. N. Tchana, J.M. Mérillon, P.F. Moundipa, B.T. Ngadjui and B.M. Abegaz, *Planta Med.*, 77: 1044–1047 (2011).
11. F. Zelefack, D. Guilet, A. Valentin, R.C. Soh Fongang, B. Kom, S. Chevalley, S.A. Ngouela, E. Tsamo, N. Fabre and M.G. Dijoux-Franca, *Greener J. Bio. Sci.*, 2 (2): 20-24 (2012).
12. R.P. Adams, *Identification of essential oil components by gas chromatography/quadrupole mass spectrometry*, 4<sup>th</sup> ed., Allured publishing, Carol Stream, IL (2007).

**S. A. Aboaba, O. Adeeko and G. Flamini**

13. T. Patharakorn, T. Arpornsuwan, N. Wetprasit, A. Promboon and S. Ratanapo, *J. Med. Pl. Res.*, 4(9): 837-843 (2010).
14. E. Stenhagen, S. Abrahamsson and F.W. McLarfferty, *Registry of Mass spectral data*. John Wiley and Sons, New York (1974).
15. Y. Massada, *Analysis of Essential Oils by Gas Chromatography and Mass spectrometry*. John Wiley and Sons, New York (1976).
16. W. Jennings and T. Shibamoto, *Qualitative analysis of flavor and fragrance volatiles by Glass Capillary Chromatography*. Academic press, New York (1980).
17. N.W. Davies, *Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicon and Carbowax 20M phases*. *J. Chromatogr., A* **503**: 1–24 (1990).
18. M. Lavy, A. Zuker, E. Leminsohn, O. Larkov, U. Ravid, A. Vainstein and D. Weiss, *Molecular breeding*, 9: 103-111 (2002).
19. J. Gertsch, M. Leenti, S. Raduner, I. Racz, J. Chen, X. Xie, K. Aktmann, M. Karsak and A. Zimmer, *PNAS*, 105(26): 9099-9104 (2008).
20. A. Bahi, S. Al Mansouri, E. Al Memari, M. Al Ameri, S.M. Nurulain and S. Ojha, *Physiol. & Behavior*, 135: 119-124 (2014).
21. R.E. Duncan, D. Lau, A. El-Sohevy and M.C. Archer, *Biochem. Pharmacol.*, 68(9): 1739-1747 (2004).
22. Y. Yao, X. Ding, Y. Jia, C. Huang, Y. Wang and Y. Xu, *Cancer Letters*, 264(1): 127-134 (2008).
23. J. Liu, Y. Zhang, J. Qu, L. Xu, K. Hou, J. Zhang, X. Qu and Y. Liu, *BMC Cancer*, 11: 183-192 (2011).

24. M. Ndungu, W. Lwande, A. Hassanali, L. Moreka and S.C. Chhabra, *Entomologia Experimentalis et Applicata*, 76(3): 217-222 (1995).
25. E. Innocent, N.K. Gikonyo and M.H.H. Nkunya, *Tanzania J. Health Res.*, 10(1): 50-54 (2008).
26. S. Schulz, S. Yildizhan and J.J. Van Loon, *J. Chem. Ecol.*, 37(4): 360-363 (2011).
27. <sup>a</sup>M.R. Tellez, K.K. Schrader and M. Kobaisy, *J. Agric Food Chem.*, 49: 5989-5992 (2001).
28. <sup>b,f,k,l</sup>N. Radulovic<sup>1</sup>, N. Dordevic<sup>1</sup>, M. Markovic and R. Palic, *Bull. Chem. Soc. Ethiop.* 24(1), 67-76 (2010).
29. <sup>d</sup>J.R. Kuate, P.H.A. Zollo, G. Lamaty, C. Menut and J.M. Basseire, *Flavor Fragr. J.*, 14:167-169 (1999).
30. <sup>e</sup>P. Weyerstah, H. Marshall, K. Thefeld and G.C. Subba, *Flavor Fragr. J.*, 13: 377-388 (1998).
31. <sup>g</sup>R.G. Binder and R.A. Faith, *J. Agric Food Chem.*, 37: 734-736 (1989).
32. <sup>h</sup>E. Campeol, G. Flamini, P.L. Goni, and I. Moreli, *J. Agric Food Chem.* 51: 1994-1999 (2003).
33. <sup>i</sup>M. Couladis, V. Tsortanidou, J. Fransisco-Ortega, A. Santos-Guerra, and C. Harvala, *Flavor Fragr. J.*, 16: 103-106 (2001).
34. <sup>j</sup>P. Weyerstahl, H. Marshall, I. Seelmann, and A. Rustaiyan, *Flavor. Fragr. J.*, 12:71-78 (1997).
35. <sup>k</sup>J.A. Pino, J. Mesa, Y. Munzo, M.P. Marti and R. Marbot., *J. Agric Food Chem.* 53: 2213-2223 (2005).

THE CHEMICAL AND MEDICINAL POTENTIALS OF THE FRUIT ESSENTIAL OIL OF  
*Chrysophyllum cainito* (INDIA STAR APPLE)

Ogunmola, O. O. Bolaji, O.S. & Sodamade, A.

Department of Chemistry

Emmanuel Alayande College of Education

OYO, NIGERIA

ABSTRACT

*Chrysophyllum cainito* is a lesser known fruit with several medicinal applications. The essential oil was obtained by hydrodistillation and was analysed by gas chromatography/mass spectrometry, GC/GC-MS. Twenty components were identified in the essential oil, the oil was characterized by a high proportion of fatty acids (69.37%) represented by pyruvic acid, 10-hendecenoic acid, E-9-tetradecenoic acid, pentadecanoic acid, hexadecanoic acid and cis-9-octadecenoic acid. The oil yield was 0.19 v/w of the wet sample and its compositional profile showed markedly qualitative and quantitative variation with essential oil from Cuba. This essential oil was seen to be active against gram-positive bacteria, *Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella typhimurium* and *Proteus mirabilis* and also gram negative bacteria, *Staphylococcus aureus* and *Streptococcus agalactiae*.

**KEYWORDS:** *Chrysophyllum cainito*, essential oil, hydrodistillation, fatty acid

RESUMO

O óleo essencial do fruto de *Chrysophyllum cainito* foi obtido por hidrodestilação e analisado com cromatografia gasosa e espectrometria de massa (GC-MS). Um total de vinte constituintes foram identificados no óleo essencial, que foi caracterizado pela presença alta de ácidos graxos (69.37%). Os componentes principais foram os ácidos piruvico, 10-hexadecenoico, E-9-tetradecenoico, pentadecenoico, hexadecenoico e cis-9-octadecenoico. O rendimento foi 0.19 volume/peso da amostra molhada e o perfil composicional foi muito diferente do óleo essencial de Cuba tanto qualitativamente quanto quantitativamente. O óleo essencial mostrou atividade antimicrobica tanto contra bactéria gram-positiva com *Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella typhimurium* e *Proteus mirabilis* quanto gram-negativa como *Staphylococcus aureus* e *Streptococcus agalactiae*.

**PALAVRAS CHAVE:** *Chrysophyllum cainito*, Óleo essencial, Hidrodestilação, Ácidos graxos

E-Mail; [ogunmolaoluranti@yahoo.com](mailto:ogunmolaoluranti@yahoo.com) or [layibolaji@gmail.com](mailto:layibolaji@gmail.com) or

[abbeyelectrochemical@hotmail.com](mailto:abbeyelectrochemical@hotmail.com)

VISIT OUR SITE: <http://www.sbjchem.he.com.br>

*Fruit Essential Oil of India Star Apple*

**INTRODUCTION**

Traditional and folklore medicine plays an important role in health care services around the globe. Plants are known to synthesize a wide range of chemical substances, many of which have been and can be of tremendous value in treatment and prevention of diseases. Man has depended on the plants and plants extract as a source of medicine, food, shelter, clothing etc. since creation [15].

Herbal drugs have been one of the oldest practiced by mankind because of their cheap availability and wide application, scientist in various parts of the country concentrate on studies of these remedies in order to use these as an alternative to expensive imported drugs [12] several pharmaceutical companies are engaged in the development of natural product drugs through the isolation of the so-called active molecule from plant extracts. It is estimated that today, plant material are present in, or have provided the models for 50% western drugs [14].

*Chrysophyllum cainito* commonly called star apple belongs to the family sapotaceae, it usually comes in two forms, either the dark purple skinned variety with red-purple pulp or the green skinned variety with clear white pulp. In Nigeria, this fruit is not commonly eaten like the African star apple fruit, *chrysophyllum albidum*. This plant is a medicinal plant which contain substances that could be used for therapeutic purposes of which are precursors for the synthesis of useful drugs [1].

In Ivory Coast, decoction of the leaves is used for the treatment of hypertension, while in Cuba, a decoction of the leaves is used as a cancer remedy, while that of the bark is used as an antitussive (cough suppressant) [9]. In Eastern part of Africa, infusion of the leaves have been used against diabetes and articular rheumatism, while the bark is considered a tonic and stimulant. In Venezuela, the fruit and the seed are used as a diuretic and the ripe fruits because of their mucilaginous character are eaten to sooth inflammation, laryngitis and pneumonia. The latex of the tree in Brazil is applied on abscesses [7]. In Nigeria, few people eat the fruit because it is not common but those that eat it do so because of the sweetness and its mucilaginous character.

*Chrysophyllum cainito* belongs to the family *sapotaceae* and their chemical and biological studies are scarce [2] and there is lack of chemical information about the genus *chrysophyllum*, but a number of reports concerning the isolation and characterization of bioactive compounds from various parts of the plant have appeared in the literature.

The volatile constituents of star apple were extracted in Cuba [10], the essential oil were analyzed by GC-MS and one hundred and four compounds were identified in the aroma concentrate of which (*E*)-2-hexanal, limonene, linalool, copaene and hexadecanoic acid were found to be the major constituents.

In another study from Florida, fresh fruit of *chrysophyllum cainito* were extracted with methanol and partitioned with hexane and ethyl acetate sequentially. Nine known polyphenolic antioxidants were identified and epicatechin is present in highest concentration [6].

*Fruit Essential Oil of India Star Apple*

The aqueous extract of the leaves was found at a dose of 20g/l to reduce the hyperglycemia of diabetic rat from 5g/l to 1.4g/l [5].

The extract of *chrysophyllum cainito* was used in goats against *Haemonchus contostus*[4] it was reported that the extract has nematocidal activity and this confirmed its use in ethnoveterinary practices and animal health management [11].

The purpose of the present investigation therefore was to extract, analyse and determine the antimicrobial activity of the essential oil of this lesser known fruit of *chrysophyllum cainito*.

## **MATERIALS AND METHODS**

### **Plant material and essential oil isolation**

Fresh fruits of *chrysophyllum cainito* (dark purple skinned variety) were collected from the campus of Ajayi Crowther University, Oyo in South West Nigeria in March, 2013. The sample was authenticated at the Forestry Research Institute of Nigeria (FRIN), Ibadan, where voucher specimens were deposited and was given voucher number FHI 109475. A sample of 700g was pulverized using mortar and pestle. The fresh fruit was subjected to hydro-distillation method using an all glass Clevenger apparatus according to procedure described by European Pharmacopoeia [18]. The oil was collected and kept in refrigerator without further treatment until GC/GC-MS analysis.

### **Gas Chromatography/GC mass spectrometric analysis**

The chemical composition of the essential oil was analysed using GC-MS technique. The mass spectrometer was SHIMADZU GCMS-QP2010 plus (Shimadzu Corporation, Japan) in the electron impact (EI) ionization mode (70ev) and HP 5MS (bonded 0.25 $\mu$ m capillary column (restek, Bellefonte, PA) Injector and detector temperature was held at 60 $^{\circ}$ C for 30 minutes then programmed to 240 $^{\circ}$ C at rate of 5 $^{\circ}$ C/min. Helium (99.99%) was the carrier gas at a flow rate of 1ml/min. Diluted samples (1/100 in hexane v/v) of 1.0ml were injected automatically, the linear velocity of the column was 36.8cm/sec, each peak was then analysed and assigned a number in the order that it was detected. The identification of the components was based on comparison of their mass spectra with those of NIST library, mass spectra database and literature.

### **Antimicrobial Activity Tests**

The antibacterial activities of the essential oil was measured against gram-negative (*Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella typhimurium* and *Proteus mirabelis* and gram-positive *Staphylococcus aureus* and *Streptococcus agalactiae* using a well diffusion method according to the National Committee for Clinical Laboratory Standard [8]. Briefly, Petri plates containing approximately 25-30ml of nutrient agar medium were swabbed using cotton applicator with a 24 hours sub-cultured bacterial strains which were prepared in dilution to match the turbidity intensity of the MacFarland Standard. Wells (6mm diameter) were punched in the agar and filled with 10 $\mu$ l of the extracts of different concentrations (1000, 100 and 10 $\mu$ gml $^{-1}$ ). The plates were incubated at 37 $^{\circ}$ C for 24 hours. The antibacterial activities were assessed by measuring the inhibition zone diameter (mm) around the well.

*Fruit Essential Oil of India Star Apple*

**RESULTS AND DISCUSSION**

Hydro distillation of the fresh fruit of *chrysophyllum cainito* produced a clear light essential oil. The oil yield is 0.19% v/w of the wet sample. The chemical components identified by GC/GC-MS are listed in Table 1. It was characterized by a high proportion of fatty acids (69.37%) represented by (IIz) –II hexadecenoic acid (24.42%), cis-9-octadecenoic acid (oleic acid) (13.04%), E-9- Tetradecenoic (12.21%), E-9-Hexadecanoic acid (9.54%), Pentadecanoic acid (3.18%), Nonanoic acid (pelargic acid) (3.18%), Pyruvic acid (2.97%) and 10-Hendecenoic acid (0.83%).

The compositional profile of the fruit oil showed markedly qualitative and quantitative variation with that from Cuba. (*E*)-2-hexanal, limonene, linalool and copaene which are part of the major constituents of the oil from Cuba were not identified from the Nigerian oil sample, but hexadecanoic acid which was reported as one of the major constituents [10] is present in the Nigerian sample (9.54%).

The oil is seen at different concentration to be active against gram-positive bacteria (*Klebsiella pneumoniae*, *Salmonella typhimurium* and *Proteus mirabilis*) and also gram-negative bacteria *Staphylococcus aureus* and *Streptococcus agalactiae* as listed in Tables 2 and 3 and this can be compared with standard antibiotics as listed in Tables 4 and 5.

*O. O. Ogunmola, O. S. Bolaji and A. Sodamide*

**Table 1. Chemical Constituents of the fruit essential oil of *chrysophyllum cainito***

S/N	COMPONENT	RI	% COMPOSITION
1	3-ethoxy-1-butane	620	0.80
2	Divinyl sulfide	650	2.97
3	2,2-Dimethylpentanal	821	0.80
4	2-pentanethiol	837	2.97
5	Heptan-3-one (ethylbutylketone)	853	0.80
6	Tert-Butylcarbinol	876	0.80
7	Pyruvic acid	919	2.97
8	Monochloromethylisopentanoate	946	0.80
9	Allyl pentanoate	974	2.97
10	Methylcyclooctane	1020	0.83
11	2(3H)-Furanone	1061	2.97
12	(4Z)-4-Methyl-4-undecene	1199	0.83
13	Nonanoic acid (pelargic acid)	1272	3.18
14	10-Hendecenoic acid (Sevinon)	1461	0.0
15	E-9-Tetradecenoic acid	1777	12.21
16	Pentadecanoic acid (pentadecyclic acid)	1869	3.18
17	Hexadecanoic acid (palmitic acid)	1968	9.54
18	(11z)-11-Hexadecenoic acid	1976	24.42
19	E-2-Octadecadecen-1-0l	2061	12.21
20	Cis-9-octadecenoic acid (oleic acid)	2175	13.04
21	Methyl(13E)-13-docosenoate	2483	0.83
	Total		99.95%

Table 2.

Zones of inhibition (mm) showing the antimicrobial activities of the essential oil at different concentrations (Gram-negative bacteria)

Organism	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Salmonella typhimorium</i>	<i>Proteus mirabilis</i>
Conc ( $\mu\text{gml}^{-1}$ )	1000 100 10	1000 100 10	1000 100 10	1000 100 10
Zone of inhibition	15 08 09	13 13 13	12 12 12	13 13 13

Keynote -- no inhibition, 6-9mm = low inhibition 10-15mm = moderate inhibition  $\geq 15\text{mm}$  = high inhibition.

Table 3.

Zones of inhibition (mm) showing the antimicrobial activities of the essential oil at different concentrations (Gram-positive bacteria).

Organism	<i>Staphyococcus aureus</i>	<i>Streptococcus agalatae</i>
Conc ( $\mu\text{gml}^{-1}$ )	1000 100 10	1000 100 10
Zone of inhibition	17 13 13	15 15 15

Keynote -- no inhibition, 6-9mm = low inhibition 10-15mm = moderate inhibition  $\geq 15\text{mm}$  = high inhibition.

*O. O. Ogunmola, O. S. Bolaji and A. Sodamade*

**Table 4.**

**Zones of inhibition (mm) showing the antimicrobial activity of antibiotics (Gram –negative bacteria)**

Organism / Antibiotics/ concentrations	<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Salmonella typhimorium</i>	<i>Proteus mirabilis</i>
Nitrofurantoin (200µg)	25	20	15	15
Ofloxacin (5µg)	20	20	22	33
Gentamicin (10µg)	24	15	18	16

Keynote --= no inhibition, 6-9mm = low inhibition 10-15mm = moderate inhibition ≥15mm = high inhibition.

**Table 5.**

**Zones of inhibition (mm) showing the antimicrobial activities of antibiotics (Gram –positive bacteria)**

Organism / Antibiotics/ concentrations	<i>Staphylococcus Aureus</i>	<i>Streptococcus agalactiae</i>
Nitrofurantoin (200µg)	28	11
Ofloxacin (5µg)	20	-
Gentamicin (10µg)	-	-

Keynote --= no inhibition, 6-9mm = low inhibition 10-15mm = moderate inhibition ≥15mm = high inhibition.

*Fruit Essential Oil of India Star Apple*

**CONCLUSION**

The results obtained in this study showed that the fruit essential oil of *chrysophyllum cainito* from South West Nigeria has a qualitative and quantitative variation from the one obtained in Cuba and this can be traced to the tropical nature of Nigeria. This study is the first in Nigeria, to the best of my knowledge.

**REFERENCES**

1. O.A. Abolaji, A.H. Adebayo and O.S. Odesanmi, (2007). Nutritional qualities of three medicinal plant parts (*xylopiya aetheopica*, *bligha sapida* and *parinar polyandra*) commonly used by pregnant women in Western part of Nigeria. *Pakistan Journal of Nutrition* 6:665-668
2. V.C. Dasilva, M.N. Lopes and V.S. Bolzani (2006). Chemical study of leaves of *chrysophyllum margination* Nova 20:3 retrieved online from <http://dx.doi.org/10.1590/150100-40422> on 07-11-2014.
3. European Pharmacopoeia (2008). retrieved from <https://www.edqm.eu>. on 08-11-2014
4. J.B. Githiori, S. Athanasiadou and S.M. Thamsborg (2006). Use of plant in novel approaches for control of gastrointestinal helminthes in livestock with emphasis on small ruminants *Veterinary Parasitology* 139:308-320
5. N. Koffi, A.K. Ernest, Marie-Solange, K. Beogreand Z.G. Noel (2009). Effect of aqueous extract of *chrysophyllum cainito* leaves on the glycemia of diabetic rabbits. *African Journal of Pharmacy and Pharmacology*. 3 (10): 500-506

O. O. Ogunmola, O. S. Bolaji and A. Sodamade

6. Luo, X., Basile, M.J. and Kennelly, E.J. (2002). Polyphenolic antioxidant from the fruits of *chrysophyllum cainito* L. (star apple). *Journal of Agric Food Chemistry* 50 (6): 1379-1382
7. J. Morton, (1987). Star Apple in fruits of Warm Climates, Julica F. Miams Morton. 408
8. National committee for clinical laboratory standards (1993). Retrieved online from [www.nchi.nlm.nih.gov/pmc](http://www.nchi.nlm.nih.gov/pmc) on 17-07-14
9. National Research Council (2008). "Star apples". Retrieved from <http://books.Napedu/openph> on 17-07-2014.
10. J. Pino, R. Marbot and A. Rosado (2002). Volatile constituents of star apple *chrysophyllum cainito* L from Cuba. *Flavour and Fragrance Journal*. 17 (5): 401 -403
11. S. Rochfort, A. J. Parker and F.B. Dunshea (2008). Plant bioactivities for ruminant health and productivity. *Phytochemistry*, 69, 299-322. Retrieved online at [www.elsevier.com/locate/photochem](http://www.elsevier.com/locate/photochem).
12. A. Sofowora (1993). *Medicinal Plant and Traditional Medicines in Africa*, Chichester John Wiley and Sons, New York. 20 – 23.
13. T.A. Tor-Anyiin, R. Sha'ato and H. O. A. Oluma (2006). Phytochemical screening and antibacterial activity of *cissampelos mucronata*. A rich *menispermaceae* extract. *Journal of Pharmacy and Bioresources* 3(2):103-106
14. F. Venter and J.A. Venter (1996). *Making the most of indigenous trees*, Briza Publications, Pretoria. 260 – 263

**SYNTHESIS AND BIOLOGICAL ACTIVITIES EVALUATION OF  
SOME NEW SPIRO 1,2,4-TRIAZOLE DERIVATIVES HAVING  
SULFONAMIDE MOIETY**

Hany M. Dalloul,<sup>\*a</sup> Khaled El-nwairy,<sup>b</sup> Ali Z. Shorafa,<sup>b</sup> Ahmed Abu Samaha<sup>c</sup>

<sup>a</sup> Department of Chemistry, Faculty of Applied Science, Alaqsa University of Gaza,  
P.O.Box 4051, Gaza 76888, Palestine. E-mail: [hmdalloul@yahoo.com](mailto:hmdalloul@yahoo.com)

<sup>b</sup> Ministry of Education & Higher Education, Beirut Street, Southern Reemal,  
Gaza 76888, Palestine.

<sup>c</sup> Department of Biology, Faculty of Applied Science, Alaqsa University of Gaza,  
P.O.Box 4051, Gaza 76888, Palestine.

**ABSTRACT**

A series of new spiro 1,2,4-triazoles **V-IXa-j** were synthesized by the reaction of appropriate amidrazones **IV** with cyclic ketones in the presence of *p*-toluene sulfonic acid as a catalyst. The structures of the synthesized compounds have been confirmed by the elemental analysis and spectroscopic data (IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR and MS). The microbial features of the synthesized compounds were studied using well-established methods from literature.

**KEYWORDS:** Amidrazone, nitrilimines, spiro 1,2,4-triazole, sulfonamide, cyclic ketone.


---

\*Corresponding author. E-mail: [hmdalloul60@yahoo.com](mailto:hmdalloul60@yahoo.com)

**GRAPHICAL ABSTRACT**

**Synthesis and Biological Activities Evaluation of Some New Spiro  
1,2,4-Triazole Derivatives Containing Sulfonamide Moiety**

**Hany M. Dalloul, Khaled El-nwairy, Ali Z. Shorafa and  
Ahmed Abu Samaha**  
Alaqa University of Gaza, Palestine



**V**, Ar = Me; **VI**, Ar = Ph; **VII**, Ar = 2-Furyl;  
**VIII**, Ar = 2-Tienyl; **IX**, Ar = 2-Naphthyl

## 1. Introduction

Amidrazones represent a class of substances with interesting biological properties. It has been established that they can exhibit antibacterial and antifungal [1-3], antitumor [4], or antituberculosis activities [5]. They were also found as effective herbicides [6], pesticides [7], and insecticides [8]. The interest in amidrazones and their derivatives stems not only from their biological relevance but also from their applications as precursors and intermediates for the synthesis of many heterocyclic compounds [9-11] or as ligands in coordination chemistry. Amidrazones synthesized from different hydrazonoyl halides are reported to react with  $\alpha$ -haloesters in the presence of triethylamine as a base under reflux afforded 1,3,5-substituted 4,5-dihydro-1,2,4-triazin-6-ones [12]. Several studies, involving the formation and investigation of biological activities of some spiroheterocyclic compounds having triazole, tetrazine thiadiazole, thiadiazines, thiazolidinone and triazine moieties [13-17]. Recently, unknown dispiroheterocycles containing triazole and tetrazine moieties have been reported by Dalloul [14-17].

Sulfonamide derivatives exhibit a range of bioactivities, including anti-angiogenic, anti-tumor, anti-inflammatory and anti-analgesic, anti-tubercular, anti-glaucoma, anti-HIV, cytotoxic, anti-microbial and anti-malarial agents [18]. Taking into account all previous commentaries of the biological activities of sulfonamides and in continuation of our study on the synthesis of biologically active heterocycles [19-21], efforts have been made to synthesize a series of new spiro heterocycles containing 1,2,4-triazole and piperidone derivatives via cyclization of amidrazones having sulfonamide moiety with cyclic ketones in anticipation of expected interesting biological activities.


## 2. Results and Discussion

The precursors of amidrazones hydrazonoyl halides **I** employed in this study were prepared according to reported literature procedures [22-25]. Treatment of the hydrazonoyl halides **I** with sodium azide in presence of tetrabutylammonium iodide at room temperature gave azidohydrazones **II** (Figure). The obtained hydrazonoyl azides **II** reacted with triphenylphosphine afforded phosphonimines **III** (Figure). It was found that the hydrolysis of compounds **III** with aqueous hydrochloric acid gave the triphenylphosphine oxide and the amidrazones **IV**, respectively (Figure). The most plausible mechanism for this acid hydrolysis is the one that involves initial protonation of nitrogen followed by the attack of oxygen on the phosphorous atom [22].

The condensation of amidrazones **IV** with cyclic ketones in refluxing dioxane in the presence of catalytic amount of *p*-toluenesulfonic acid produced spiro 4,5-dihydro-1H-1,2,4-triazole derivatives containing sulfonamide moiety **V-IXa-j** (Figure) in good yields. The reaction progress was monitored by TLC to find out the completion of the synthesis.

### 2.2. Spectral data analysis of compounds **IV** and **V-IXa-j**.

The structure of the newly synthesized amidrazones **IV** was elucidated on the basis of their spectroscopic data and elemental analyses. The electron impact (EI) mass spectra of these compounds **4a-j** displayed the correct molecular ions ( $M^+$ ) in accordance with the suggested structures. The IR spectra exhibit typical stretching absorption bands of C=O of conjugate ketone, anilide and ester groups at about 1725-1650  $\text{cm}^{-1}$ , NH's in the region of 3470-3220  $\text{cm}^{-1}$  and 1150, 1060  $\text{cm}^{-1}$  attributed to  $\text{SO}_2$  of sulfonamide group. The  $^1\text{H}$  NMR spectra of these amidrazones DMSO- $d_6$  show three characteristic signals, as singlet at 12.6-12.4 ppm ( $\text{SO}_2\text{NH}$ ), singlet of the NNH group in the region of 6.3-6.0 ppm and singlet of  $\text{NH}_2$  near 4.9-5.0 ppm. In addition the characteristic singlet signal of the CONH group in compounds containing anilide group was observed at about 10.8-9.8 ppm and the expected proton signals of the aromatic rings in the range of 8.4-6.9 ppm. The  $^{13}\text{C}$  NMR spectra exhibit characteristic signals for the Ar-C=O carbon at about 192-159 ppm and for the C=N moiety at about 143-141 ppm.



**Figure.** Synthetic pathway for the preparation of compounds **IV** and **V-IXa-j**.

The structural assignment of the prepared spiro 1,2,4-triazoles **V-IXa-j** was based on elemental analysis and spectral data. The electron impact (EI) mass spectra of these compounds **4a-j** displayed the correct molecular ions ( $M^+$ ) in accordance with the suggested structures. Physical properties, molecular ion peaks and microanalysis are presented in experimental section. The IR spectra of these compounds revealed the presence of NH band of dihydrotriazole ring resonated near  $3370-3350\text{ cm}^{-1}$ . In addition to the bands of characteristic functional groups. The  $^1\text{H}$  NMR spectra of products **V-IXa-j** in  $\text{DMSO-d}_6$  display a characteristic singlet in the region of 5.6-5.5 ppm due to NH proton of dihydrotriazole ring. Also, the spectra exhibit a characteristic singlet at 12.7-12.6 ppm due to  $\text{SO}_2\text{NH}$  proton. The  $^{13}\text{C}$  NMR spectra display characteristic signals of the suggested structures. The signal at 90-95 ppm, which is attributed to C-5 (spiro carbon) of dihydrotriazole ring is of special significance. This is similar to reported values of spiro carbons flanked by two nitrogens in five-membered heterocycles [23,26]. The spectral data of the obtained compounds **IVa-j** are summarized in the experimental section.

## **2.2. Antimicrobial activity**

The standard nutrient agar disc diffusion method<sup>35,36</sup> was followed to determine the activity of the synthesized compounds against the sensitive organisms *Eutero cocci*, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella spp*, *Proteus spp*, as bacterial strains and two species of fungi, namely *Aspergillus niger*, *Candida albicans*. The compounds tested at a concentration of  $1\text{ mg mL}^{-1}$  in N,N-dimethyl formamide (DMF) solution, and measuring the average diameter of the inhibition zone in mm. The results showed that all the tested compounds exhibited a marked degree of activity against bacteria and fungi compared with well-known antibacterial and antifungal substances such as tetracycline and fluconazole. According to NCCLS [27], zones of inhibition for tetracycline and fluconazole  $< 14\text{ mm}$  were considered resistant, between 15 and 18 mm were considered weakly sensitive and  $> 19\text{ mm}$  were considered sensitive. Also, the results showed the degree of inhibition varied with the tested compounds (Table). The thiazolyl and pyrimidinyl moieties generally led to dramatic improvements in activity against both bacteria and fungi. The present study can lead medicinal chemists to design and synthesize similar compounds with enhanced biological potency in future.

**Table.** Antimicrobial screening results of the tested compounds

Cpd. No.	Diameter of the inhibition zone in mm*						
	Antibacterial activity					Antifungal activity	
	<i>Eutero cocci</i>	<i>Escherichia coli</i>	<i>Staphylo aureus</i>	<i>Klebsiella spp</i>	<i>Proteus spp</i>	<i>Candida albicans</i>	<i>Aspergillus niger</i>
<b>Va</b>	16	17	17	13	16	15	14
<b>Vb</b>	13	18	15	11	10	17	18
<b>Vc</b>	19	15	11	14	16	18	16
<b>Vd</b>	18	16	17	18	19	16	12
<b>VIb</b>	16	19	16	19	11	19	11
<b>VIIa</b>	13	12	14	16	17	16	19
<b>VIIb</b>	19	15	11	14	16	18	16
<b>VIIIa</b>	18	16	17	18	19	16	12
<b>VIIIb</b>	16	19	16	19	11	19	11
<b>IXa</b>	13	12	14	16	17	16	19
DMF	–	–	–	–	–	–	–

\*Calculated as average of three values.

### 3. EXPERIMENTAL SECTION

#### 3.1. Reagents and Instrumentation

Melting points were determined on an A. Krüss Melting Point Meter equipped with a thermometer and are uncorrected. The IR spectra were measured as potassium bromide pellets using a Satellite 3000 Mid infrared spectrophotometer. The <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on a Bruker AM 300 MHz spectrometer at room temperature in DMSO-d<sub>6</sub> solution using tetramethylsilane (TMS) as internal reference. Chemical shifts were recorded as δ values in parts per millions (ppm) downfield from internal TMS. Electron impact (EI) mass spectra were run on a Shimadzu GCMS-QP1000 EX spectrometer at 70 eV. All compounds were analyzed satisfactorily for C, H and N. The amidrazones **IV** were prepared according to literature procedures [22-25]. Sodium azide, triphenylphosphine and tetrabutylammonium iodide were obtained from Fluka Chemie Company, Switzerland. Sulfathiazole, sulfadiazine, dioxane, tetrahydrofuran (THF) and triethylamine were purchased from Avocado Research Chemicals, England, and used without further purification.

#### 3.2. Synthesis of spiro 4,5-dihydro-1,2,4-triazole derivatives V-IXa-j.

To a stirred solution of the appropriate amidrazone **IV** (10 mmol) and the respective cycloalkanone (20 mmol) in dioxane (50 mL), the catalytic amount of *p*-toluenesulfonic acid (0.1 g) was added. The reaction mixture was refluxed to the completion (monitoring the reaction progress by TLC). The excess of the solvent was evaporated

and the residue was triturated with methanol or ethanol. The resulting crude solid product was collected and recrystallized from ethanol. The following compounds were prepared using this method:

**3-Acetyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene**

**(Va):** Yield: 65%; m.p.: 197-199 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3382, 3355 (NH's), 1695 ( $\text{CH}_3\text{-C=O}$ ), 1622 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.70 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 4H, Ar-H), 8.71 (*d*, 1H, thiazole), 6.62 (*d*, 1H, thiazole), 5.65 (*s*, 1H, NH triazole ring), 2.45 (*s*, 3H,  $\text{COCH}_3$ ), 2.37-1.60 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  192.3 (C=O), 169.1-116.5 (Ar-C, C=N and thiazole-C), 92.8 (spiro-C), 34.9, 24.9 (cyclopentane-C), 26.5 ( $\text{CH}_3$ ); MS:  $m/z$  405 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{17}\text{H}_{19}\text{N}_5\text{O}_3\text{S}_2$  (405.50): C, 50.35; H, 4.72; N, 17.27; Found: C, 50.57; H, 4.60; N, 17.15.

**3-Acetyl-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene**

**(Vb):** Yield: 71%; m.p.: 216-218 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3374, 3350 (NH's), 1693 ( $\text{CH}_3\text{-C=O}$ ), 1626 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.74 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 4H, Ar-H), 8.41 (*d*, 2H, pyrimidine), 6.83 (*t*, 1H, pyrimidine), 5.63 (*s*, 1H, NH triazole ring), 2.55 (*s*, 3H,  $\text{COCH}_3$ ), 2.37-1.60 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  192.1 (C=O), 168.19-110.6 (Ar-C, C=N and pyrimidine-C), 92.6 (spiro-C), 34.8, 24.6 (cyclopentane-C), 26.6 ( $\text{CH}_3$ ); MS:  $m/z$  400 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_3\text{S}$  (400.46): C, 53.99; H, 5.03; N, 20.99; Found: C, 54.23; H, 4.92; N, 21.10.

**3-Acetyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene**

**(Vc):** Yield: 66%; m.p.: 204-206 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3369, 3357 (NH's), 1695 ( $\text{CH}_3\text{-C=O}$ ), 1624 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.76 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 4H, Ar-H), 8.71 (*d*, 1H, thiazole), 6.65 (*d*, 1H, thiazole), 5.65 (*s*, 1H, NH triazole ring), 2.54 (*s*, 3H,  $\text{COCH}_3$ ), 2.07-1.16 (*m*, 10H, cyclohexane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  192.4 (C=O), 168.4-116.2 (Ar-C, C=N and thiazole-C), 92.1 (spiro-C), 35.9, 24.8, 23.2 (cyclohexane-C), 26.5 ( $\text{CH}_3$ ); MS:  $m/z$  419 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{18}\text{H}_{21}\text{N}_5\text{O}_3\text{S}_2$  (419.53): C, 61.53; H, 5.05; N, 16.69; Found: C, 61.70; H, 4.96; N, 16.55.

**3-Acetyl-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene**

**(Vd):** Yield: 65%; m.p.: 192-194 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3377, 3357 (NH's), 1693 ( $\text{CH}_3\text{-C=O}$ ), 1628 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.71 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 4H, Ar-H), 8.40 (*d*, 2H, pyrimidine), 6.82 (*t*, 1H, pyrimidine), 5.61 (*s*, 1H, NH triazole ring), 2.55 (*s*, 3H,  $\text{COCH}_3$ ), 1.98-1.16 (*m*, 10H, cyclohexane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  191.9 (C=O), 168.25-110.4 (Ar-C, C=N and pyrimidine-C), 91.8 (spiro-C), 35.8, 24.6, 23.1 (cyclohexane-C), 26.4 ( $\text{CH}_3$ ); MS:  $m/z$  414 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{19}\text{H}_{22}\text{N}_6\text{O}_2\text{S}$  (414.49): C, 55.06; H, 5.35; N, 20.28; Found: C, 54.78; H, 5.50; N, 20.39.

**3-Acetyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.6]undec-2-ene**

**(Ve):** Yield: 67%; m.p.: 178-180 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3375, 3358 (NH's), 1695 ( $\text{CH}_3\text{-C=O}$ ), 1625 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.76 (*s*, 1H, NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.70 (*d*, 1H, thiazole), 6.64 (*d*, 1H, thiazole), 5.63 (*s*, 1H, NH triazole ring), 2.54 (*s*, 3H,  $\text{COCH}_3$ ), 2.35-1.42 (*m*, 12H, cycloheptane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  192.4 (C=O), 168.2-116.5 (Ar-C, C=N and thiazole-C), 91.6 (spiro-C), 39.4, 28.1, 22.3 (cycloheptane-C), 26.5 ( $\text{CH}_3$ ); MS:  $m/z$  433 [ $\text{M}^+$ ]; Anal. Calcd. for

C<sub>19</sub>H<sub>23</sub>N<sub>5</sub>O<sub>3</sub>S<sub>2</sub> (433.55): C, 52.64; H, 5.35; N, 16.15; Found: C, 52.45; H, 5.24; N, 16.03.

**3-Acetyl-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.6]undec-2-ene (Vf):** Yield: 64%; m.p.: 188-190 °C; IR (KBr)  $\nu_{\max}$ : cm<sup>-1</sup> 3373, 3365 (NH's), 1695 (CH<sub>3</sub>-C=O), 1620 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.69 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.36 (*d*, 2H, pyrimidine), 6.81 (*t*, 1H, pyrimidine), 5.62 (*s*, 1H, NH triazole ring), 2.56 (*s*, 3H, COCH<sub>3</sub>), 2.34-1.40 (*m*, 12H, cycloheptane); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  191.8 (C=O), 168.25-110.4 (Ar-C, C=N and pyrimidine-C), 39.1, 28.3, 22.5 (cycloheptane-C), 91.8 (spiro-C), 26.4 (CH<sub>3</sub>); MS: *m/z* 428 [M<sup>+</sup>]; Anal. Calcd. for C<sub>20</sub>H<sub>24</sub>N<sub>6</sub>O<sub>3</sub>S (428.52): C, 56.06; H, 5.65; N, 19.61; Found: C, 55.86; H, 5.53; N, 19.47.

**3-Acetyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.7]dodec-2-ene (Vg):** Yield: 70%; m.p.: 166-168 °C; IR (KBr)  $\nu_{\max}$ : cm<sup>-1</sup> 3373, 3361 (NH), 1696 (CH<sub>3</sub>-C=O), 1627 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.71 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.72 (*d*, 1H, thiazole), 6.61 (*d*, 1H, thiazole), 5.65 (*s*, 1H, NH triazole ring), 2.55 (*s*, 3H, COCH<sub>3</sub>), 2.42-1.24 (*m*, 14H, cyclooctane); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  192.3 (C=O), 168.05-116.5 (Ar-C, C=N and thiazole-C), 90.9 (spiro-C), 41.9, 36.5, 27.2, 21.8 (cyclooctane-C), 26.5 (CH<sub>3</sub>); MS: *m/z* 447 [M<sup>+</sup>]; Anal. Calcd. for C<sub>20</sub>H<sub>25</sub>N<sub>5</sub>O<sub>3</sub>S<sub>2</sub> (447.58): C, 53.67; H, 5.63; N, 15.65; Found: C, 53.45; H, 5.75; N, 15.55.

**3-Acetyl-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.7]dodec-2-ene (Vh):** Yield: 65%; m.p.: 169-171 °C; IR (KBr)  $\nu_{\max}$ : cm<sup>-1</sup> 3366, 3360 (NH's), 1692 (CH<sub>3</sub>-C=O), 1626 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.76 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.40 (*d*, 2H, pyrimidine), 6.81 (*t*, 1H, pyrimidine), 5.66 (*s*, 1H, NH triazole ring), 2.55 (*s*, 3H, COCH<sub>3</sub>), 2.42-1.24 (*m*, 14H, cyclooctane); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  192.2 (C=O), 168.1-110.2 (Ar-C, C=N and pyrimidine-C), 91.8 (spiro-C), 41.8, 36.3, 27.1, 21.6 (cyclooctane-C), 26.6 (CH<sub>3</sub>); MS: *m/z* 442 [M<sup>+</sup>]; Anal. Calcd. for C<sub>21</sub>H<sub>26</sub>N<sub>6</sub>O<sub>3</sub>S (442.54): C, 57.00; H, 5.92; N, 18.99; Found: C, 56.85; H, 6.05; N, 18.87.

**3-Acetyl-8-methyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene (Vi):** Yield: 71%; m.p.: 186-188 °C; IR (KBr)  $\nu_{\max}$ : cm<sup>-1</sup> 3374, 3355 (NH's), 1693 (CH<sub>3</sub>-C=O), 1622 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.70 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.71 (*d*, 1H, thiazole), 6.65 (*d*, 1H, thiazole), 5.64 (*s*, 1H, NH triazole ring), 2.54 (*s*, 3H, COCH<sub>3</sub>), 2.07-1.14 (*m*, 8H, cyclohexane), 0.98 (3H, *s*, CH<sub>3</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  189.9 (C=O), 168.2-116.5 (Ar-C, C=N and thiazole-C), 34.9, 27.8, 22.4 (cyclohexane-C), 90.9 (spiro-C), 31.4 (CH<sub>3</sub> at cyclohexane ring), 26.5 (CH<sub>3</sub>); MS: *m/z* 433 [M<sup>+</sup>]; Anal. Calcd. for C<sub>19</sub>H<sub>23</sub>N<sub>5</sub>O<sub>3</sub>S<sub>2</sub> (433.55): C, 52.64; H, 5.35; N, 16.15; Found: C, 52.85; H, 5.20; N, 16.28.

**3-Acetyl-8-tert-butyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene (Vj):** Yield: 69%; m.p.: 199-201 °C; IR (KBr)  $\nu_{\max}$ : cm<sup>-1</sup> 3370, 3353, (NH's), 1695 (CH<sub>3</sub>-C=O) 1622 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.69 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 4H, Ar-H), 8.68 (*d*, 1H, thiazole), 6.63 (*d*, 1H, thiazole), 5.57 (*s*, 1H, NH triazole ring), 2.57 (*s*, 3H, COCH<sub>3</sub>), 2.05-1.13 (*m*, 8H, cyclohexane), 0.92 (9H, *s*, (CH<sub>3</sub>)<sub>3</sub>C); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  189.9 (C=O), 168.2-116.5 (Ar-C, C=N and thiazole-C), 90.8 (spiro-C), 35.2, 27.9, 22.6 (cyclohexane-C), 46.9, 35.7, 32.4, 24.0 (*tert*-butyl carbons), 26.4 (CH<sub>3</sub>); MS: *m/z* 470 [M<sup>+</sup>]; Anal. Calcd. for C<sub>23</sub>H<sub>30</sub>N<sub>6</sub>O<sub>3</sub>S (470.60): C, 58.70; H, 6.43; N, 17.86; Found: C, 58.55; H, 6.31; N, 17.95.

**3-Benzoyl-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (VIa):** Yield: 66%; m.p.: 219-221 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3368, 3345 (NH's), 1675 (Ph-C=O), 1596 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.56 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 9H, Ar-H), 8.74 (*d*, 1H, thiazole), 6.60 (*d*, 1H, thiazole), 5.63 (*s*, 1H, NH triazole ring), 89.8 (spiro-C), 2.37-1.60 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  185.2 (C=O), 167.9-116.2 (Ar-C, C=N and thiazole-C), 34.6, 24.7 (cyclopentane-C); MS:  $m/z$  467 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{22}\text{H}_{21}\text{N}_5\text{O}_3\text{S}_2$  (467.57): C, 56.51; H, 4.53; N, 14.98; Found: C, 56.25; H, 4.40; N, 15.10.

**3-Benzoyl-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (VIb):** Yield: 67%; m.p.: 236-238 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3371, 3347 (NH's), 1676 (Ph-C=O), 1597 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.58 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 9H, Ar-H), 8.44 (*d*, 2H, pyrimidine), 6.81 (*t*, 1H, pyrimidine), 5.58 (*s*, 1H, NH triazole ring), 2.37-1.60 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  185.5 (C=O), 168.1-110.2 (Ar-C, C=N and pyrimidine-C), 89.9 (spiro-C), 34.8, 24.7 (cyclopentane-C); MS:  $m/z$  462 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{23}\text{H}_{22}\text{N}_6\text{O}_3\text{S}$  (462.53): C, 59.73; H, 4.79; N, 18.17; Found: C, 59.55; H, 4.95; N, 18.05.

**3-(2-Furoyl)-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (VIIa):** Yield: 70%; m.p.: 197-199 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3375, 3355 (NH's), 1665 (Ar-C=O), 1616 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.63 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 7.82-7.06 (*m*, 7H, Ar-H), 8.66 (*d*, 1H, thiazole), 6.58 (*d*, 1H, thiazole), 5.65 (*s*, 1H, NH triazole ring), 2.37-1.60 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  173.6 (C=O), 168.2-116.4 (Ar-C, C=N and thiazole-C), 90.6 (spiro-C), 34.7, 24.6 (cyclopentane-C); MS:  $m/z$  457 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{20}\text{H}_{19}\text{N}_5\text{O}_4\text{S}_2$  (457.53): C, 52.50; H, 4.19; N, 15.31; Found: C, 52.67; H, 4.06; N, 15.43.

**3-(2-Furoyl)-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene (VIIc):** Yield: 71%; m.p.: 226-228 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3377, 3352 (NH's), 1666 (Ar-C=O), 1615 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.64 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 7H, Ar-H), 8.67 (*d*, 1H, thiazole), 6.56 (*d*, 1H, thiazole), 5.60 (*s*, 1H, NH triazole ring), 2.05-1.12 (*m*, 10H, cyclohexane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  173.5 (C=O), 168.2-116.5 (Ar-C, C=N and thiazole-C), 90.4 (spiro-C), 36.2, 25.1, 23.2 (cyclohexane-C); MS:  $m/z$  471 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{21}\text{H}_{21}\text{N}_5\text{O}_4\text{S}_2$  (471.56): C, 53.49; H, 4.49; N, 14.85; Found: C, 53.35; H, 4.55; N, 14.73.

**3-(2-Thenoyl)-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (VIIIa):** Yield: 72%; m.p.: 218-220 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3365, 3345 (NH's), 1665 (Ar-C=O), 1612 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.66 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 7H, Ar-H), 8.70 (*d*, 1H, thiazole), 6.63 (*d*, 1H, thiazole), 5.63 (*s*, 1H, NH triazole ring), 2.35-1.58 (*m*, 8H, cyclopentane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  174.6 (C=O), 168.1-116.2 (Ar-C, C=N and thiazole-C), 88.8 (spiro-C), 34.8, 24.9 (cyclopentane-C); MS:  $m/z$  473 [ $\text{M}^+$ ]; Anal. Calcd. for  $\text{C}_{20}\text{H}_{19}\text{N}_5\text{O}_3\text{S}_2$  (473.60): C, 50.72; H, 4.04; N, 14.79; Found: C, 50.90; H, 3.90; N, 14.65.

**3-(2-Thenoyl)-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.5]dec-2-ene (VIIId):** Yield: 73%; m.p.: 198-200 °C; IR (KBr)  $\nu_{\max}$ :  $\text{cm}^{-1}$  3365, 3346 (NH's), 1660 (Ar-C=O), 1610 (C=N), 1346, 1175 ( $\text{SO}_2$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  11.59 (*s*, 1H,  $\text{SO}_2\text{NH}$ ), 8.27-7.06 (*m*, 7H, Ar-H), 8.38 (*d*, 2H, pyrimidine), 6.81 (*t*, 1H, pyrimidine), 5.61 (*s*, 1H, NH triazole ring), 2.04-1.13 (*m*, 10H, cyclohexane);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  174.8 (C=O), 168.4-110.5 (Ar-C, C=N and pyrimidine-C), 88.9 (spiro-C),

36.1, 25.3, 23.1 (cyclohexane-C); MS:  $m/z$  482 [ $M^+$ ]; Anal. Calcd. for  $C_{22}H_{22}N_6O_3S_2$  (482.59): C, 54.76; H, 4.60; N, 17.41; Found: C, 53.60; H, 4.05; N, 19.35.

**3-(2-Naphthoyl)-1-[4-(thiazol-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (IXa):** Yield: 61%; m.p.: 212-213 °C; IR (KBr)  $\nu_{max}$ :  $cm^{-1}$  3369, 3350 (NH), 1606 (Ar-C=O), 1608 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.66 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 11H, Ar-H), 8.76 (*d*, 1H, thiazole), 6.66 (*d*, 1H, thiazole), 5.53 (*s*, 1H, NH triazole ring), 2.37-1.60 (*m*, 8H, cyclopentane); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  185.4 (C=O), 168.0-116.2 (Ar-C, C=N and thiazole-C), 89.8 (spiro-C), 34.9, 24.9 (cyclopentane-C); MS:  $m/z$  517 [ $M^+$ ]; Anal. Calcd. for  $C_{26}H_{23}N_5O_3S_2$  (517.63): C, 60.33; H, 4.48; N, 13.53; Found: C, 60.15; H, 4.60; N, 13.65.

**3-(2-Naphthoyl)-1-[4-(pyrimidin-2-yl-sulfonyl)phenyl]-1,2,4-triazaspiro[4.4]non-2-ene (IXb):** Yield: 62%; m.p.: 165-167 °C; IR (KBr)  $\nu_{max}$ :  $cm^{-1}$  3367, 3355 (NH's), 1655 (Ar-C=O), 1605 (C=N), 1346, 1175 (SO<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  11.67 (*s*, 1H, SO<sub>2</sub>NH), 8.27-7.06 (*m*, 11H, Ar-H), 8.46 (*d*, 2H, pyrimidine), 6.85 (*t*, 1H, pyrimidine), 5.55 (*s*, 1H, NH triazole ring), 2.34-1.51 (*m*, 8H, cyclopentane); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  185.6 (C=O), 168.2-110.4 (Ar-C, C=N and pyrimidine-C), 89.7 (spiro-C), 34.8, 24.7 (cyclopentane-C); MS:  $m/z$  512 [ $M^+$ ]; Anal. Calcd. for  $C_{27}H_{24}N_6O_3S$  (512.59): C, 63.27; H, 4.72; N, 16.40; Found: C, 63.40; H, 4.62; N, 16.28.

#### 4. CONCLUSION

New series of novel functionalized spiro1,2,4-triazols **V-IXa-j** bearing sulfonamide moiety were synthesized upon the treatment of amidrazones **Va-j** with cyclic ketones in refluxing dioxane and evaluated for their in vitro antibacterial, and antifungal activities. From the screening results, it found to possess various antimicrobial activities towards all the microorganisms tested. The results confirm that, the antimicrobial activity is strongly dependent on the nature of the substituents on triazole ring.

#### 5. ACKNOWLEDGMENTS

The authors are great thankful to the Qatar Charity for the financial support of this research through Ibhath grant (GCC-07-06).

#### 6. REFERENCES

- [1] Mamolo, M. G.; Vio, L.; Banfi, E.; Predominato, M.; Fabris, C.; Asaro, F. *IL Farmaco* **47**, 1055 (1992).
- [2] Mamolo, M. G.; Vio, L.; Banfi, E.; Predominato, M.; Fabris, C.; Asaro, F. *IL Farmaco* **48**, 529 (1993).
- [3] Ranft, D.; Seyfarth, T.; Schaper, K. J.; Lehwerk-Yvetot, G.; Bruhn, C.; Buge, A. *Arch Pharm Pharm Med Chem* **332**, 427 (1992).
- [4] Modzelewska-Banachiewicz, B.; Matysiak, J.; Niewiadomy, A. *Eur J Med Chem* **2001**, 36, 75.

- [5] Abdel-Jalil, R.; J.; Al Momani, E.; Hamad, M.; Voelter, W.; Mubarak, M. S.; Smith, B. H.; Peters, D. G. *Monatsh Chem* 2010, 141, 251.
- [6] Dean, G. S.; Rhodes, S. G.; Coad, M.; Whelan, A. O.; Wheeler, P.; Villareal-Ramos, B.; Mead, E.; Johnson, L.; Clifford, D. J.; Hewinson, R. G.; Vordermeier, H. M. *Tuberculosis* 2008, 88, 586.
- [7] Hackmann, J. T. US Patent 2 758 050, 1956.
- [8] Haldeman, R. G.; Morin, L. T.; Matsuda, K. US Patent 3 073013, 1963.
- [9] Kuhn, D. G.; Furch, J. A.; Hunt, D. A.; Asselin, M.; Baffic, S. P.; Diehl, R. E.; Miller, T. P.; Palmer, Y. L.; Treacy, M. F.; Trotto, S. H. *ACS Symp Ser* 1998, 686, 185.
- [10] Neilson, D. G.; Roger, R.; Heatlie, J. W. M.; Newlands, L. R. *Chem Rev* 1970, 70, 151.
- [11] Aly, A. A.; Nour-El-Din, A. M. *Arkivoc* **i**, 153 (2008).
- [12] Dalloul, H. M.; Boyle, P. H. *Turk. J. Chem.* **2007**, *31*, 141-147.
- [13] Borad, M.A.; Bhoia, N.; Prajapatia, N. P.; Patela, H. D. *Synth. Commun.* **2014**, *44*(7), 897-922; *44*(8), 1-15.
- [14] Shawali, A. S.; Abdelhamid, A. O. *Curr.Org. Chem.* **2014**, (17) 2673-2689.
- [15] Dalloul, H. M. *Arkivoc* **2008**, (xiv) 234-241.
- [16] Dalloul, H. M.; Abu Samaha, A. S. *J. Serb. Chem. Soc.* **2010**, *75*, 1473.
- [17] Dalloul, H. M. *South. Braz. J. Chem. Soc.* **2014**, *22*, 17-26.
- [18] Mok, B. L. PhD, Chemistry Department, University College London, UK, **2008**.
- [19] Dalloul, H. M. *Can. Chem. Trans.*, **2015**, *3*(1), 108-117;
- [20] Dalloul, H. M. *J. Chem.*, 1-7 (2015).
- [21] Dalloul, H. M. *Open J. Org. Chem.*, **3**(2), 12-17 (2015).
- [22] Shawali, A. S.; Hassaneen, H. M.; Fahmy, A. A.; Abunada, N. M.; Mousa, H. A. *Phosphorus, Sulfur. and Silicon*, **53**, 259-265 (1990).
- [23] Drutkowski, G.; Donner, C.; Schulze, I.; Frohberg, P.; *Tetrahedron*, **58**, 5317-5326 (2002).
- [24] A.S. Ferwanah, A.M. Awadallah, E.A. El-Sawi and H.M. Dalloul, *Synth. Commun.* **33**, 1245-1253 (2003).
- [25] H.M. Dalloul and P.H. Boyle, *Heterocycl. Commun.* **9**, 507 (2003).
- [26] Dalloul, H. M., *Chem. Heterocycl. Comp.*, **40**(11), 1402-1407 (2004).
- [27] Clinical and Laboratory Standards Institute, "Methods for dilution antimicrobial susceptibility tests for bacteria grow aerobically", Approved Standard M7-A4, Clinical and Laboratory Standards Institute, Wayne, Pa, USA, 2005.

**KINETICS OF REACTION BETWEEN MALACHITE GREEN AND HYDROXYL ION  
IN THE PRESENCE OF REDUCING SUGARS.**

**<sup>1</sup>Latona, Dayo Felix and <sup>2</sup>Dada, Adewumi Oluwasogo**

<sup>1</sup>Department of Chemical Sciences, Osun State University, Osogbo, Nigeria.

<sup>2</sup>Industrial Chemistry Unit, Department of Physical Sciences, Landmark University,  
Omu-Aran, Nigeria.

Corresponding author's e-mail: [d1latona@yahoo.com](mailto:d1latona@yahoo.com) or [dayo.latona@uniosun.edu.ng](mailto:dayo.latona@uniosun.edu.ng)

**Abstract**

The reaction was studied via pseudo first order kinetics using uv-1800 Shimadzu spectrophotometer with a thermostated cell compartment and interfaced with a computer . The reaction showed first order with respect to malachite green and sugar and hydroxyl ion concentrations. However, the reaction was independent of ionic strength and showed no dependence on salt effect, indicating an inner sphere mechanism for the reaction. There was no polymerization of the reaction mixture with acrylonitrile, indicating absence of radicals in the course of the reaction. Michaelis-Menten plot indicated the presence of a reaction intermediate in the rate determining step. The activation parameters of the reaction have been calculated and products were elucidated by FTIR spectroscopy. The stoichiometry of the reaction is 1:1. A mechanism consistent with the above facts has been suggested.

**Keywords:** Malachite green(MG), sodium hydroxide, Glucose, Fructose, Xylose.

DOI: 10.48141/SBJCHEM.v23.n23.2015.61\_revista2015.pdf

## Introduction

Malachite green is a triphenylmethane dye which has wide range of industrial applications, most especially in aquaculture production because of its antibacterial, antifungal and antiparasitic properties(1) . However, they are toxic to human as they have been confirmed to be mutagenic and carcinogenic (2). Owing to its hazardous effect, several researches have been reported on the best ways to remove it in water or industrial effluents mainly by biosorption(3,4,5,6). It has been established that wood fibre of phoenix tree( *Firmiana Simplex*) is an effective adsorbent for malachite green(7).

Fading of malachite green(MG) in the presence of hydroxyl ion has been attributed to the hydroxide ion attack on the central C atom of the planar ring system of malachite green, thereby destroying its conjugation orientation. It has been found that sugars such as dextrose, lactose and sucrose increases the rate of fading of dyes in general (8). Hitherto, there has been no report on the kinetics and mechanism of alkaline malachite green fading in sugars. However, few or scanty investigations have been reported on the kinetics of alkaline malachite green fading in alcohols and in the presence of surfactants(9,10).

Hence the need to study the effect of some reducing sugars on the alkaline fading of malachite green with the view to determining the mechanism via a vis the mechanism earlier proposed for the oxidation of sugars in sodium hydroxide medium(11).

## Materials and methods

Analar grade malachite green, NaOH, KOH, glucose, fructose and xylose were used and the solutions were prepared in doubly distilled water.

The reaction products were characterized by FTIR and the reaction showed no presence of radicals as there were no polymerization on addition of acrylonitrile to the reaction mixture.

## Spectral Measurement

$\lambda_{\text{max}}$  of dilute solution of malachite green was obtained at 617 nm using a uv-1800 Shimadzu spectrophotometer.

## Kinetic Measurements

The reactions were performed under pseudo-first order conditions by maintaining a large excess (x10 or greater) of sugar over malachite green. The kinetic data were obtained by monitoring change in absorbance of malachite green at absorption maximum as a function of

time with a UV-1800 Shimadzu spectrophotometer connected to a thermostated cell compartment and interfaced with a computer. Appropriate quantities of the solution of sugar, potassium nitrate, sodium hydroxide and malachite green were measured into the cuvette. Consequently, the reaction was kick started after adding requisite volume of the malachite green solution ( All stock solutions were kept in the water bath for 30 minutes before the kinetic runs). The kinetic data were obtained via pseudo- first order condition with the concentration of the sugar in large excess compared with the oxidant concentration. The pseudo- first order rate constant ( $k_{obs}$ ) were calculated.


## **Results and Discussion**

### Stoichiometry and product analysis

The stoichiometry of the reaction was determined by spectroscopic titration. Absorbances of solutions containing various concentrations of sugar within the range  $1.00 \times 10^{-2}$  to  $6.00 \times 10^{-2}$  and a constant  $[MG]_0$  of  $1.00 \times 10^{-3}$  and constant  $[NaOH]_0$  of  $6.00 \times 10^{-2}$  and  $[KNO_3]_0$  of 0.25M were measured at 710nm after the reaction had gone to completion. The stoichiometry was evaluated from the plot of absorbance vs [sugar] curve and was found to be 1:1.

### **Effect of [MG] on the observed rate constant**

A plot of  $\ln k_{obs}$  vs  $\ln[MG]$  gave a slope equal to 1, indicating a first order dependence. The reaction showed an increase in rate constant with increase in [MG] as shown in fig.1.

Fig. 1: Plot of  $k_{\text{obs}}$  vs [MG]

(● Fructose ■ Glucose ▲ Xylose)

### Effect of [sugar] on the observed rate constant

As the concentration of sugars increases, there is increase in the rate constants as shown in table 1. The plot of  $\ln k_{\text{obs}}$  vs  $\ln[\text{sugar}]$  shows that the reaction increases with increase in sugar concentrations, indicating that the complex formation took place between sugar and malachite green and the slope of this plot is unity, implying a first order dependence on [sugar]. The plot of  $1/k_{\text{obs}}$  vs  $1/[\text{s}]$  gave an intercept, revealing the presence of an intermediate complex as shown in fig.2. Second order rate constants are  $0.152 \text{ M}^{-1}\text{S}^{-1}$  (Glucose),  $0.138 \text{ M}^{-1}\text{S}^{-1}$  (Fructose),  $0.125 \text{ M}^{-1}\text{S}^{-1}$  (Xylose).

**Table 1: Effect of [sugar] on reaction rate**

$10^2[S]/M$	$10^3k_{obs}/s^{-1}$		
	Glucose	Fructose	Xylose
1.00	1.81	2.10	1.73
2.00	3.60	3.40	2.81
3.00	5.12	4.75	3.62
4.00	6.84	6.02	4.80
6.00	8.10	7.41	6.51

[MG]  $1.00 \times 10^{-3}M$  [OH<sup>-</sup>]  $6.00 \times 10^{-2}M$  I= 0.25M T=298K

**Effect of [OH<sup>-</sup>] on the observed rate constant**

Increase in [OH<sup>-</sup>] resulted into increased reaction rate as shown in table 2 . The slope of a plot of  $\ln k_{obs}$  vs  $\ln[OH^-]$  also showed first order dependence.


**Table 2: Effect of [OH<sup>-</sup>] on reaction rate**

$10^2[OH^-]/M$	$10^3k_{obs}/s^{-1}$		
	Glucose	Fructose	Xylose
1.00	1.98	1.81	2.20
2.00	3.14	2.30	3.00
3.00	4.22	3.82	4.21
4.00	5.05	4.85	5.30
5.00	6.70	6.25	6.00
6.00	8.10	7.41	6.91

[MG]  $1.00 \times 10^{-3}M$  [S]  $2.00 \times 10^{-2}M$  I= 0.25M T= 298K

**Effect of ionic strength on the observed rate constant**


The reaction rate was independent of the ionic strength of the solution. A plot of  $\ln k_{obs}$  vs  $I^{1/2}$  gave a straight line with slope equal to zero as shown in fig.3, which implies the presence of a neutral molecule in the rate determining step.

Fig.2: Plot of  $1/k_{obs}$  vs  $1/[S]$ 

(● Fructose ■ Glucose ▲ Xylose)

#### Salt effect on the observed rate constant

The addition of KCl and NaNO<sub>3</sub> salts did not affect the reaction rate as shown in fig.4. This reveals an inner sphere mechanism for the reaction.

Fig. 3: Plot of  $\ln k_{\text{obs}}$  vs  $I^{1/2}$ 

### Effect of temperature and determination of activated parameters

The effect of temperature on the observed rate constants were determined by varying the temperature within 298K- 308K and concentrations of all the reactants were constant:  $[\text{MG}] 1.00 \times 10^{-3}$ ,  $[\text{sugar}] 2.00 \times 10^{-2}\text{M}$ ,  $[\text{OH}^-] 6.00 \times 10^{-2}\text{M}$  and  $I= 0.25\text{M}$ . The observed rates constants at different temperatures were determined and the values of activation parameters were calculated using the below relationship and shown in table 3. The result showed that the rate of reaction

increases with increase in temperature as this is evident by the high negative values of entropies of activation. Moreover, nearly constant values of Gibbs' free energies of activation suggests a similar mechanism for all the sugars.

**Table 3: Activation parameters**

Sugar	Ea(kJmol <sup>-1</sup> )	ΔH <sup>#</sup> (kJmol <sup>-1</sup> )	ΔS <sup>#</sup> (kJK <sup>-1</sup> mol <sup>-1</sup> )	ΔG <sup>#</sup> (kJmol <sup>-1</sup> )
Glucose	23.52	21.04	-0.182	75.28
Fructose	25.12	22.64	-0.177	75.38
Xylose	31.34	28.86	-0.156	75.35

[MG] 1.00 x 10<sup>-3</sup>M [sugar] 2.00 10<sup>-2</sup>M [OH<sup>-</sup>] 6.00 X 10<sup>-2</sup>M I= 0.25M T=298K

$$\log k = \log A - \frac{E_a}{2.303RT}$$

$$\ln\left(\frac{k}{T}\right) = \frac{-\Delta H^\#}{RT} + \ln\left(\frac{k'}{h}\right) + \left(\frac{\Delta S^\#}{R}\right)$$

$$\ln\left(\frac{k'}{h}\right) = 23.76$$

$$\Delta G^\# = \Delta H^\# - T\Delta S^\#$$

k = Rate constant

T= Temperature

ΔH<sup>#</sup>= Enthalpy of activation


ΔS<sup>#</sup>= Entropy of activation

ΔG<sup>#</sup> = Free Gibb's energy of activation

R= Molar gas constant

k' = Boltzmann's constant

h= Plank's constant




FTIR Spectrum of the product.

### Mechanism and Rate Law:

The non-dependence of the oxidation reaction on ionic strength indicates the presence of a neutral molecule in the rate determining step. Same values of  $\Delta G^\ddagger$  indicates same mechanism for the reaction and negative values of  $\Delta S^\ddagger$  revealed entropy decrease upon achieving the transition state, which often indicate an associative mechanism and it shows that the reaction occurred between ions of similar charges. The FTIR spectrum of the product showed very broad  $-\text{OH}$  stretching at  $3400\text{--}2400\text{ cm}^{-1}$ ,  $\text{C}=\text{O}$  stretching at  $1730\text{--}1700\text{ cm}^{-1}$  and  $\text{C}-\text{O}$  stretching between  $1320\text{--}1210\text{ cm}^{-1}$  which can be attributed to the presence of carboxylic acid. Michaelis-Menten plot showed the presence of an intermediate complex. Relatively low values of  $\Delta H^\ddagger$  revealed the absence of high energy free radicals.

Considering the above results, the following reaction mechanism is hereby proposed. The alkaline fading of malachite green in the presence of sugars proceeds via the conversion of sugars to their corresponding enediols which of course is a fast step. Then the enediol reacts with malachite green via an outer sphere mechanism to give a complex (slow step) which in turn decomposes into products.



$$\frac{d[E]}{dt} = k_1[S][OH^-] - k_{-1}[E] - k_2[E][MG] \quad (2)$$

$$0 = k_1[S][OH^-] - k_{-1}[E] - k_2[E][MG]$$

$$k_1[S][OH^-] = k_{-1}[E] + k_2[E][MG]$$

$$[E] = \frac{k_1[S][OH^-]}{k_{-1} + k_2[MG]} \quad (3)$$

From equations (1) and (3)

$$\frac{-d[MG]}{dt} = \frac{k_2 k_1 [S][OH^-][MG]}{k_{-1} + k_2[MG]} \quad (4)$$

*Limiting condition*

If  $k_{-1}$  is far greater than  $k_2[MG]$ . Then eq. 4 becomes

$$\frac{-d[MG]}{dt} = K_1 k_2 [S][OH^-][MG]$$

## Conclusion

The research work showed that the mechanism of alkaline malachite green fading in reducing sugar is similar to that of the oxidation of sugars in alkaline medium. The reaction product as shown by the FTIR spectrum revealed the presence of carboxylic acid. This implies that sugars are been oxidized to their corresponding acids by malachite green and malachite green was decolorized via reduction process. However, in the absence of sugar, hydroxyl ion decolorizes malachite green by attacking the central C atom of the planar ring system of malachite green, thereby destroying its conjugation orientation which could have resulted in the production of carbinol.

## Acknowledgement

We appreciate the technical support given by the technical staff of the central science laboratory, Obafemi Awolowo University, Ile-Ife, Nigeria.

## Reference

1. M.M. El Jamal , M.AEI Hajj Hassan and L.M.A. Fayoumi (2011). Kinetic study of the discoloration of triphenyl methane dyes in function of  $P^H$ , salt effect. Journal of the Uni. Chem. Tech. and Metallurgy,46,4, 395-400.
2. S. Srivastava , R. Sinha and D. Roy (2004). Toxicological effects of malachite green. Aquatic Toxicology, 66, 319-329.
3. R. Deokar and A.Sabale (2014). Biosorption of Methyhlene Blue and Malachite Green from Binary Solution onto Ulva Lactuca. Int. J. Curr. Microbiol. App.Sci. 3(5), 295-304.
4. J. Singh and G. Kaur (2013). Freundlich, Langmuir adsorption isotherms and kinetics of the removal of malachite green from aqueous solutions using agricultural waste rice straw. Int. Journal of Environ. Sci. vol. 4, No 3,250-255.
5. M. Hema and S. Arivoli (2008). Adsorption kinetics and thermodynamics of malachite green dye unto activated low cost carbon. J. Appl. Sci. Environ. Manage. Vol.12(1) 43-51.

6. M. Santhi , P.E.Kumar and B. Muralidharan (2015). Removal of malachite green dye by adsorption onto activated carbon-MnO<sub>2</sub>-Nanocomposite- kinetic study and equilibrium Isotherm analyses. IOSR Journal of Applied chemistry vol. 8, Issue 4, 33-41.
7. D. Zhang and X.Pan (2009). Removal of malachite green from waste by Firmiana Simplex wood fibre. Electronic Journal of Biotechnology vol. 12 No 4, 1-10.
8. R.Kuramoto, L. Lachman and J. Cooper (1958). A study of the effect of certain pharmaceutical materials on color stability. Journal of Amer. Pharma. Ass., Vol. XLVII, No 3, 175-180.
9. A. Raducan, A. Olteanu , M. Pius and D.Dancea .(2008). Influence of surfactants on the fading of malachite green. Central European Journal of chemistry 6(1), 89-92.
10. B. Samiey and A.R. Toosi (2009). Kinetic study of malachite green fading in the presence of TX-100, DTAB and SDS. Bull Korean Chem. Soc. Vol. 30, No 9, 2051-2056.
11. D.F. Latona (2016). Kinetics and mechanism of sugar oxidation by KMnO<sub>4</sub> in NaOH medium. The Pacific Journal of Science and Technology, Vol. 17 No 1, 177-182.

CHEMICAL CONSTITUENTS AND PHYTOCHEMICAL STUDIES OF  
EPAZOTE (*Chenopodium ambrosioides*) LINN GROWN  
IN NORTH CENTRAL NIGERIA

Larayetan R.A<sup>1\*</sup>; AdegbeA, A<sup>1</sup>; Owonikoko A.D.<sup>2</sup>

<sup>1</sup> Department of Chemistry, Kogi State University  
Anyigba, Kogi State, NIGERIA

<sup>2</sup> Department of Chemistry, Emmanuel Alayande College of Education,  
Oyo State, NIGERIA

ABSTRACT

The essential oil from the leaves of *C.ambrosioides* of North Central Nigeria was obtained by hydro-distillation using Clevenger apparatus. The oil was analysed by a combination of GC and GCMS. Nineteen components amounting to 98.37% of the total oil were identified. The main constituents of the North Central Nigeria grown *C.ambrosioides* are 2-Carene (17.80%), 2-bornene (14.79%), p-Cymene (12.93%),  $\alpha$ -Terpinene (8.98%), Cyclohexene-4-methyl-3-(1-methylethylidene) (7.94%),  $\alpha$ -Terpinolene (7.90%),  $\gamma$ -Terpinene (6.94%). The oil was found to be a yellow liquid, the obtained yield is 1.35%w/w based on the dry weight. The preliminary phytochemical investigation showed that *C.ambrosioides* methanolic leaf extract contains some secondary metabolites such as Flavonoids, Terpenoids, Steroids, Alkaloids and Saponins. The result justified the use of *C.ambrosioides* in treating various infectious diseases.

**KEYWORDS:** Volatile constituents, Phytochemical screening, Epazote, hydro distillation, *Chenopodium ambrosioides*.

RESUMO

O óleo essencial de folhas de *C. ambrosioides* da parte Norte-Central da Nigéria foi obtido por hidrodestilação usando o aparelho de Clevenger. O óleo foi analisado com uma combinação de cromatografia gasosa e espectrometria de massa. Foram identificados 19 componentes correspondendo à 98,37% do total. Os constituintes principais de *C. ambrosioides* foram careno (17,80%), 2-borneno (14,79%), p-cimeno (12,93%), alfa-terpineno (7,90%), ciclohexeno-4-metil-3(1-metiletilideno) (7,94%), alfa-terpinoleno (7,90%) e gama-terpineno (6,94%). O óleo é um líquido amarelo e corresponde à 1,35% por peso do peso seco. Estudos preliminares fitoquímicos indicaram que as folhas contêm flavonoides, terpenoides, steróis, alcalóides e saponinas e justificam o uso de *C. ambrosioides* no tratamento de doenças infecciosas.

**PALAVRAS CHAVE:** *Chenopodium ambrosioides*, Óleo essencial, Hidrodestilação, Estudos fitoquímicos, Constituintes voláteis

\*Corresponding Author: Larayetan Rotimi. A. Department of Chemistry, Kogi State University, Anyigba, Nigeria. Email Address: [timlarayetan@gmail.com](mailto:timlarayetan@gmail.com). Phone number: +2348030510909

## **INTRODUCTION**

Higher and aromatic plants have been used traditionally in folk medicine as well as to extend the shelf life of foods, showing inhibition against bacteria, fungi and yeast. Biologically active compounds from natural sources have always been of great interest to scientists [1]. The essential oils known as volatile oils evaporate in contact with air and possess a pleasant fragrance. Chemically essential oils are very complex. They are found in many different species of plants of various families. All aromatic plants contain essential oils. Generally the oils are secreted in the oil glands [2]

According to WHO, about three quarter of the world population relies upon traditional remedies [mainly herbs] for the health care of its people. In fact plants are the oldest friends of mankind. They not only provide food and shelter but also serve the humanity to cure different ailments [3].

The plant family Chenopodiaceae is a large family comprising about 102 genera and 1400 species [4]. The genus *Chenopodium* include varieties of weedy herbs (more than 200 species) native to Europe, Asia and both North and South America [5]. *Chenopodium ambrosioides* Linn also called (Epazote, Mexican tea, Wormseed e.t.c) is a widespread species native to tropical America [6]. It is annual or short-lived perennial herbs that have been used for centuries as condiment, traditional purgative for intestinal worms and many other medicinal purposes [7-9]. The leaves and seeds are green when fresh. The plant is sometimes cultivated principally for its medicinal use in West Africa as purgative, for treating sores, edema and flavouring purposes [10]. They are employed for different ailments in Africa. The importance of *Chenopodium* species is due to their wide variety of medicinal properties [11].

*C.ambrosioides* is called Ewe-imi, Asin and Arunpale in some parts of Nigeria [12]. The whole plant leaves are used as anthelmintics, emollient and to treat rheumatism and tumour in parts of the country [13]. It has also been employed by empirical herbalist and healers against intestinal parasites (especially small tapeworms and roundworms) throughout Latin America; as well as West Indies [14]. As a result of its effect against intestinal parasites, traditional healers in North central Nigeria administer about 2-3 gm of the dried powdered leaves of *C.ambrosioides* mixed with pap to treat hookworm infection and hookworm inflammatory diseases, the decoction is also used in this region as wash for various skin diseases, to treat fever and as a fumigant against mosquitoes.

The biological activity of *C.ambrosioides* has been shown to affect viruses [15], bacteria [16], fungi [17], nematodes [18], and insects [19]. Furthermore the plant is traditionally and widely used as antiparasitic, anti-inflammatory and antibiotics which efficacy has been scientifically proven [20].

Chemical composition of the essential oil of *C.ambrosioides* from different parts of the world has been widely studied such as from Brazil [21], Cuba [22], Mexico [23], Cameroon [24], Nigeria [25], Rwanda [26], China [27,28], and India [29,30]. Ascaridole and Isoascaridole that were found in the Brazil grown *C.ambrosioides* [31] were absent in the North Central Nigeria grown *C.ambrosioides*, also  $\alpha$ -Terpinene that constituted the major component in the Nigeria grown *C.ambrosioides* by [Gbolade et al,32] was just (8.98%) in the North Central Nigeria grown *C.ambrosioides*, 2-carene was found to be the major constituent.

Phytochemical are bioactive compounds found in plants they are non-nutritive compound (secondary metabolites) that contributed to flavour, colour [33, 34]. They are needed by plant for purpose such as disease, pathogen defence and control. The aim of this study was to identify the volatile constituents of the leaf oil of *C.ambrosioides* grown in North Central Nigeria by using modern GC/MS machine [SHIMADZU GCMS-QP2010 PLUS], as well as to find out the phytochemical components in the methanol extracts of *C.ambrosioides* grown in this region.

## **MATERIALS AND METHODS**

### **Collection of plant leaves and identification**

Fresh matured leaves of the plant were collected in May 2014 from a village in Anyigba of Kogi State, North Central Nigeria. They were identified by Dr Aina of the Department of Botany, Kogi State University, Anyigba. Voucher specimens are deposited in the herbarium of the Faculty of Biological Sciences, Kogi State University, Anyigba, Nigeria

### **Isolation of the volatile oil**

The method employed for the extraction of the volatile oil of the leaf of *Chenopodium ambrosioides* was hydro distillation method according to European Pharmacopoeia (2008). Oils were collected and kept in the refrigerator without further treatment before GC-MS analysis.

### **Preparation of plant extracts of methanol**

The collected fresh matured leaves of the plant of *C.ambrosioides* were air dried usually at room temperature in a well aerated room; the air dried leaves were grinded into coarse powdered using a Thomas Willey machine. The aqueous extract was prepared by soaking 200g

of the dried powdered samples in 400ml of methanol for 24 hrs. The extracts were filtered using Whatman filtered paper No 42 (125mm).

### **Screening of Phytochemical Components**

To identify the phytochemical derivatives in the methanolic extract, standard phytochemicals screening was performed [35-37]. Alkaloid test was performed by Meyer's tests, flavonoid and tannins by ferric chloride test, terpenoid and sterols by Salkowski's test, saponin by frothing test and anthraquinones by Borntrager's test.

### **Identification Test**

The following tests were carried out so as to detect active chemical constituents such as alkaloids, tannins, flavonoids, anthraquinones, steroids, terpenoids and saponin.

#### **Test for Alkaloids**

1cm<sup>3</sup> of 1%HCl was added to 3cm<sup>3</sup> of the extract in a test tube. The extract was treated with a few drops of Meyer's reagent. A creamy white precipitate was observed indicating the presence of alkaloid.

#### **Test for Tannins**

About 0.5g of the dried powdered samples was boiled in 20ml of water in a test tube and then filtered. A few drops of 0.1% ferric chloride was added and observed for brownish green or blue back colouration. There was no brownish green or blue colouration observed.

#### **Test for Saponin**

About 2g of the powdered samples was boiled in 20ml of distilled water in a water bath and filtered. 10ml of the filtrate was mixed with 5ml of distilled water and shaken vigorously, then observed for the formation of emulsion.

#### **Test for Flavonoids**

About 0.5g of the extract was boiled with 5.0ml of distilled water and then filtered. To 2.0ml of this filtrate; a few drops of 10% ferric chloride solution were added. A greenish blue or violet colouration indicated the presence of a phenolic hydroxyl group.

### **Test for Terpenoids**

5ml of the extract was mixed in 2ml of chloroform, and concentrated H<sub>2</sub>SO<sub>4</sub> (3ml) was carefully added to form a layer. A reddish brown colouration of the inter face was formed to show positive results for the presence of terpenoids.

### **Test for Sterols**

A few milligrams of the plant extract were dissolved in 2ml of chloroform and then 2ml of concentrated H<sub>2</sub>SO<sub>4</sub> was added from the side of the test tube. The test tube was shaken for a few minutes. Red colour development in the chloroform layer indicated the presence of sterols.

### **Test for Anthraquinones**

An aliquot of 0.5g of the extract was boiled with 10ml of H<sub>2</sub>SO<sub>4</sub> and filtered while hot. The filtrate was shaken with 5ml of chloroform. The chloroform layer was pipette into another test tube and 1ml of dilute ammonia was added. The resulting solution was observed for colour change. There was no colour change.

## **2.2 Gas chromatography/mass spectrometry analysis**

The chemical composition of the essential oil was analysed using GC/MS technique. The mass spectrometer was SHIMADZU GCMS-QP2010 Plus (Shimadzu Corporation, Japan) in the electron impact (EI) ionization mode (70eV) and HP-5MS (bonded 0.25 $\mu$ m) Capillary column (restek, Bellefonte, PA). Injector and Detector temperature were set at 250°C. The oven temperature was held at 60°C for 30minutes, then programmed to 240°C at rate of samples (1/100 in hexane v/v) of 1.0ml were injected automatically. The linear velocity of the column was 36.8cm/sec, each peak was then analysed and assigned a number in the order that it was detected. The identification of the components was based on comparison of their mass spectra with those of NIST library mass spectra database and literature.

**Table 1. Volatile Constituents of the Leaf Oil of *Chenopodium Ambrosioides*.**

S/N	RI	COMPONENT	% COMPOSTION
1	932	2-Bornene	14.79
2	948	IR- $\alpha$ -pinene	1.82
3	998	$\alpha$ -Terpinene	8.98
4	1001	2- Carene	17.8
5	1010	3,7,7-trimethyl-1,3,5-cycloheptatriene.	7.94
6	1023	4-methyl-3-(1-methylidene)-1-cyclo hexane	4.31
7	1042	<i>p</i> -cymene	12.93
8	1052	$\alpha$ -Terpinolene	7.90
9	1094	Nona-3,5-diene-2-ol.	1.88
10	1119	<i>m</i> -Xylene	4.61
11	1143	$\alpha$ -Terpineol	0.50
12	1179	$\gamma$ -Terpinene	6.94
13	1195	Decyn-2-ol	1.94
14	1248	1,2:4,5:9,10-Triepoxydecane	1.88
15	1763	Z-10-pentadecen-1-ol	0.57
16	2007	9-Octadecenal	0.64
17	2085	16-Octadecenoic acid, methyl ester	1.44
18	2175	Oleic acid	1.14
19	2682	15-Tetracosenoic acid	0.36
TOTAL IDENTIFIED			98.37%

**Note; RI Means Retention Index**

## RESULTS AND DISCUSSION

The hydro distillation of the leaf oil of *C.ambrosioides* yielded 1.35% w/w essential oil. Nineteen components amounting to 98.37% were identified in the leaf oil. The identified components, their retention index and percentage composition of each component is given in table 1 above. Most of the components being monoterpenes and oxygenated monoterpenes. The major components found in the essential oil of *C.ambrosioides* were 2-Carene (17.8%), 2-Bornene (14.79%), *p*-Cymene (12.93%),  $\alpha$ -Terpinene (8.98%), Cyclohexene-4-methyl-3-(1-methylethylidene) (7.94%),  $\alpha$ -Terpinolene (7.90%) and  $\gamma$ -Terpinene (6.94%).

Constituents present in significant amount are *m*-Xylene (4.61%), 3, 7, 7-Trimethyl-1, 3, 5-cycloheptatriene (4.01%), Decyn-2-ol (1.94%), Nona-3, 5-dien-2-ol, IR- $\alpha$ -Pinene (1.82%) *e.t.c*. It is apparent that the main compounds characterizing the leaf oil are qualitatively and quantitatively different. It is also worth mentioning that compounds like *m*-Xylene, nona-3,5-dien-2-ol, decyn-2-ol, 3,7,7-trimethyl-1,3,5-cycloheptatriene and 2-bornene which are detected in the North Central Region of Nigeria samples have not been reported previously as part of the constituents of the volatile oil of *C.ambrosioides*.

The composition of the essential oil of *C.ambrosioides* has been the subject of several studies and data from literature shows that it has no constancy neither with respect to the

*R. A. Larayetan, A. A. Adegbe and A. D. Owonikoko*

relative humidity, irradiance, photoperiod and the method of extraction, location, plant cultivation techniques, soil structure and climate heavily influence the composition and quality of essential oil. [38]. These factors have been used to ascertain why exact specification of the component of essential oils is not acceptable.

The percentage constituents obtained for this sample shows that there is a marked difference in the earlier report given on *C.ambrosioides* leaf by different authors [39, 40 & 32]. The report shows that *C.ambrosioides* leaf was composed of  $\alpha$ -terpinene as the major constituent. However, the major component of the North Central Region of Nigeria *C.ambrosioides* oil was found to be 2-carene with 17.8 percentage. Ascaridole was not detected in the North Central Nigeria oil of *C.ambrosioides*.

The phytochemical screening of the methanolic extract was carried out in order to analyse the presence of secondary metabolites such as flavonoids, alkaloids, terpenoids, sterols e.t.c] by utilizing standard methods [35-37]. Table 2 shows the results of the preliminary phytochemicals analysis. This study has revealed the presences of phytochemicals considered as active medicinal chemical constituents. Important medicinal phytochemicals such as terpenoid, flavonoid, alkaloid, sterols, were present in *C.ambrosioides* leaves [Table: 2.] Plants rich in terpenoid have been reported to have anti inflammatory, antimalaria, antiviral, inhibition of cholesterol synthesis and antibacterial [41], while those rich in alkaloid are used in medicine to reduce headache and fever. These are attributed for antibacterial and analgesic properties [42]. Epidemiologic studies recommended that coronary heart diseases are opposed by dietary flavonoids. [42].

**Table 2. Phytochemical Constituents of the Leaf of *Chenopodium Ambrosioides*.**

Phytochemicals	Test	Me
Alkaloids	Meyer's Test	+
Tannins	Ferric Chloride Test	-
Saponin	Frothing Test	+
Flavonoids	Ferric Chloride test	+
Terpenoids	Salkowski Test	+
Anthraquinones	Borntrager's Test	-
Sterols	Salkowski Test	+

**Key:** (+) means Present; (-) means absent; Me means Methanolic Extract

**CONCLUSION**

The results obtained in this study showed that *C.ambrosioides* possess essential oil in the leaves of the plant and that their oil compositions were quantitatively and qualitatively different. This study represents the first to the best of my knowledge analysis of the leaves volatile constituents of *C.ambrosioides* by using a modern GCMS machine [SHIMADZU GCMS QP-2010 PLUS]. The phytochemical analysis showed the presence of effective biological compounds like alkaloids, steroids, flavonoids saponin and terpenoid in *C.ambrosioides* leaves thus providing knowledge of the phytochemical metabolites. Observation drawn from this experiment shows clearly that the leaves of *C.ambrosioides* contain phytochemical constituents. However the present investigation showed that the studied plant is a potential good source of traditional medicine and because it's a medicinal plant, it has commercial interest in both research institutes and pharmaceuticals company for the manufacture of new drugs in the treating of various diseases.

**REFERENCES**

1. K Perumal and S Ignacimuthu (2000). Antibacterial activity of some medicinal plants from Eastern Ghats, South India. *Solai Bull. Ethnopharmacol*, 72:39-41.
2. RTYousef andGGTawil (1980). Antimicrobial activity of Vol. Oils. *Die Pharmazie*, 35:698-701.
3. AHGilani and Atta-ur-Rahman (2005). Trend in Ethnopharmacology. *J Ethnopharmacol*. 100(1-2): 43-49.
4. CN.Marie In garden of Hawaii (1965). Bishop Museum Press.331.
5. BDSmith Eastern North America as an independent of plant domestication proceeds. *Nat Head Sci USA*. 103:12223-12228.
6. PAMunz(1975). A California Flora and Supplement. University of California Press Pp369.
7. MJRuffa, GFerraro, MLWagner, MLCalcagno, RHCampos andLCavallaro (2002). Cytotoxic effect of Argentine medicinal plant extracts on human hepatocellular carcinoma cell line. *J. Ethnopharm*; 79: 335-339.
8. RFFlavia, VBNascimento, SPGustavo, VPaulo,CGMMarcia, ASLucilene, SAAna Paula, SBBElizabeth andNMGRosane (2005). Ascitic and solid Ehrlich tumor inhibition by *Chenopodium ambrosioides* L. Treatment. Published by Elsevier Inc.

*R. A. Larayetan, A. A. Adegbe and A.D. Owonikoko*

9. AM Jamali L. Kouhila, AIT. Mohammed, JT. Jaouhari A Idlimam and N Abdenouri(2006). Sorption Isotherms of *Chenopodium ambrosioides* leaves at three temperatures. *J.FoodEngin* 72:77-84.
10. HM Burkill (1985). *The Useful Plants of West Tropical Africa (family A-D)*. Royal Botanic Gardens, London, Kew. Pp366.
11. LA Tapondjou,C Adler, H Bouda and DA Fontem(2002). Efficacy of powder and Essential Oil of *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six stored product beetles. *J.Stored Prod. Res*; 30:395
12. TA Taye (2013). An overview of the anti-cancer properties of some plants used in traditional medicine in Nigeria. *Int Res. J. Of Biochemistry and Bioinformatics*. Vol 3(1). Pp7-14.
13. AA Sowemimo,FA Fakoya, I Awopetu, OR Omobuwajo, SA Adesanya(2007). Toxicity and mutagenic activity of some selected Nigeria Plants. *J. Ethnopharmacol*.
14. MB Quinlan, RJQuinlan,JM Nonlan(2002).Ethno physiology and herbal treatment of intestinal worms in Dominica, West Indies. *J.Ethnopharmacol*. 80: Pp75-83.
15. HM Verma andVK Baranwal (1983). Antiviral activity and the physical properties of the leaf extract of *Chenopodium ambrosioides*. *L Proc. Indian Acad. Sci*, 92: 461-465.
- 16.JBegum,M Yusuf,JJ Chowdhury and MA Wahab (1993). Studies of essential oils for their antibacterial and antifungal properties. Part1. Preliminary Screening of 35 Essential Oils. Bangladesh. *J.Sc. Ind. Res*. 28:15-24.
17. NKDubey and N Kishore (1987). Fungi toxicity of some higher plants and synergistic activity of their oils. *Trop. Sci*; 27:23-27.
18. R Garcia-Espinosa (1980). *Chenopodium ambrosioides*. Plantacon use potencial en el combate de nematodes fitoparasitos. *Agrociencia*, Pp21-24.
- 19.GSPeterson, MA Kandil MD Abdallah and AA Farag (1989). Isolation and characterization of biologically-active compounds from some plant extracts. *Pest. Sci*; 25:343-353.
20. WSPereira, BP Ribeiro , AIP Sousa , CPB Serrai, NS Mattar, TS Fortes, AS Reis, LA Silva, ESB Barroqueiro (2010). Evaluation of subchronic toxicity of oral treatment with *Chenopodium ambrosioides* in mice. *J.Ethnopharmacol*.127:602-605. Doi.1016.
21. CM Jardim,GN Jham, OD Dhingra and MM Freire(2008) *J Chem Ecol.*, 34, 1213.
- 22.JAPino R, Marbot andIM Real (2003). *J Essent Oil Res.*, 15(3), 213.
- 23.LSagrero-Nieves and JP Bartley (1995). *J Essent Oil Res.*, (7) 221-223.
24. MSGChekem, PK Lunga, JDD, Tamokou,JR Kuate, P Tane, G Vilaremand M Cerny (2010). *Pharmaceuticals* (3), 2900-2909.
25. MSOwolabi, L Lajide, MO Oladimeji,WNSetzer, MC Palazzo, RA Olowu and A Ogundajo (2009). *Nat Prod Commun*, 4(7), 989-992.
26. AMuharyimana,JC Chalchat and RP Garry (1998). *J Essent Oil Res.*, (10), 690-692.

*Chemical Constituents and Phytochemical Studies of Epazote*

28. SSChu, JF Hu and ZL Liu (2011). *Pest Manag Sci.*, (67), 714.
29. D Gupta, R Charles, VK Mehra, SN Garg and S Kumar (2002). *J Essent Oil Res*; (14), 93
30. HPSingh, DR Batish, RK Kohli, S Mittal and S Yadav (2008). *Chem Nat Compds*; (44), 378.
31. SA Rafaela, AMG Marcos, RAF Magda, ALS Luiz and PR Karina (2014). *Chemical Composition of Essential Oil from Chenopodium ambrosioides*. *L. Rev. Bras. Farm.* 95 (3): 855-866.
32. AAGbolade, V Tira-Picos and JMF Noguera (2010). *Chemical constituents of Chenopodium ambrosioides. Varantheleminticum herb essential oil from Nigeria*. *Chem. Nat. Comp.* 46(4): 654-655.
33. A Pelter, R Warren, N Hameed, NU Khan, M Ilyas, W Tahman (1970). *Biflavonoid pigments from Thuja Orientalis (cupressaceae)*. *Phytochemistry*, (9): 1897-1898.
34. TJohn (1996). *Phytochemicals as evolutionary mediators of human nutritional Physiology* *International Journal of Pharmacognosy*, vol. 34, no. 5, Pp327-337.
35. WC Craig (1999). *Health promoting properties of common herbs*. *American journal of Clinical Nutrition*. Vol. 70, no.3 Pp 491-499.
36. WC Evans (2002). *Trease and Evans Pharmacognosy*, 15th Edition. W.B Saunders Company Ltd, London. Pp137-139, 230-240.
37. JBHarborne (1973). *Phytochemical methods: A guide to modern techniques of plant analysis*, 13th Edition. Chapman and Hall Ltd. London. Pp 5-15.
38. LPanizzi, GFlamini, PL Coni and I Morelli (1993). *J Ethnopharmacol.*
39. PAOnocha, O Ekundayo, T Eramo, and I Laasko (1999). *Essential oil constituents of C.ambrosioides L. Leaves from Nigeria*. *J. Essent Oil Res.* 11(2):220-222.
40. AA Kasali, O Ekundayo, C Paul, WA Konig, AD Eshilokun, B Ige (2006) *1,2:3,4-diepoxy-p-menthane and 1,4-epoxy-p-menth-2-ene: rare monoterpenoids from the essential oil of Chenopodium ambrosioides L. Varambrosioides leaves*. *J. Essent. Oil Res.* 18(1):13-15.
41. A Sofowora (1993). *Medicinal Plant and Traditional Medicine in Africa*; Spectrum Book Limited, Ibadan. Pp1-12, 101-108. Trease, Evans WC (1997). *Textbook of Pharmacognosy*, 14<sup>th</sup> Edition. Pp50:150.
42. SB Mahato, S Sen (1997). *Advances in triterpenoid research, 1990-1994*. *Phytochemistry* 44: 1185-1236.

VOLATILE OIL CONSTITUENTS OF THE LEAVES AND WOODS OF *Cedrela odorata*  
AND *Dalbergia latifolia* FROM SOUTH WEST NIGERIA.Ugah, T. C<sup>1</sup>, Aboaba, S. A<sup>1,\*</sup> and Ekundayo, O<sup>1</sup><sup>1</sup> Department of Chemistry, University of Ibadan

Ibadan, Nigeria

## ABSTRACT

The essential oils from the leaves and woods of *Cedrela odorata* and *Dalbergia latifolia* were obtained by hydrodistillation and characterized by Gas Chromatography-Mass Spectrometry (GC-MS) method. A total of 10, 39, 72, and 6 compounds representing 97.63%, 99.16%, 95.13% and 99.83% were identified in *Cedrela odorata* leaf, wood, *Dalbergia latifolia* leaf and wood respectively. The major components in *Cedrela odorata* leaf are  $\alpha$ -copaene (4.40%),  $\alpha$ -santalene (4.57%), cubenol (8.71%),  $\beta$ -elemene (23.06%) and (-)-spathulenol (42.49%) while the wood was dominated by  $\gamma$ -eudesmol (8.84%), z-nerolidol (9.23%),  $\beta$ -bisabolol (10.95%),  $\alpha$ -curcumene (12.31%) and  $\alpha$ -cedrene (17.56%). The leaf essential oil of *Dalbergia latifolia* had in its composition majorly  $\alpha$ -bergamotene (4.23%), (-)-spathulenol (4.49%),  $\beta$ -caryophyllene oxide (5.04%),  $\alpha$ -selinene (4.89%), 1-heptariacotanol (5.71%) and heptacosane (6.31%) while the wood contained mainly methylcyclohexane (4.93%), hexanal (6.43%), m-xylene (16.71%) and p-xylene (58.8%).

**Key words:** Volatile oil, Sesquiterpenes, (-)-spathulenol,  $\alpha$ -bergamotene,  $\beta$ -elemene

## RESUMO

Os óleos essenciais de folhas e madeira de *Cedrela odorata* e *Dalbergia latifolia* foram obtidos através de hidrodestilação e caracterizados com cromatografia gasosa e espectrometria de massa (GC-MS). Um total de 10, 39, 72 e 6 compostos representando 97.63%, 99.16%, 95.13% e 99.83% foram identificados em folhas de madeira de *Cedrela odorata* e folhas de madeira de *Dalbergia latifolia*, respectivamente. Os componentes majoritários de folhas de *Cedrela odorata* são  $\alpha$ -copaeno (4.40%),  $\alpha$ -santaleno (4.57%), cubenol (8.71%),  $\beta$ -elemeno (23.06%) e (-)-espatulenol (42.49%). Na madeira prevaleceram  $\gamma$ -eudesmol (8.84%), z-nerolidol (9.23%),  $\beta$ -bisabolol (10.95%),  $\alpha$ -curcumeno (12.31%) e  $\alpha$ -cedreno (17.56%). O óleo essencial de folhas de *Dalbergia latifolia* consistia principalmente de  $\alpha$ -bergamoteno (4.23%), (-)-espatulenol (4.49%), óxido de  $\beta$ -carofileno (5.04%),  $\alpha$ -selineno (4.89%), 1-heptarioctanol (5.71%) e heptacosano (6.31%). Na madeira prevaleceram metilciclohexano (4.93%), hexanal (6.43%), m-xileno (16.71%) e p-xileno (58.8%).

**PALAVRAS CHAVE:** Óleos voláteis, sesquiterpenos,  $\alpha$ -bergamoteno,  $\beta$ -elemeno, (-)-espatulenol

---

\* Author to whom all correspondence should be directed, e-mail: [saboaba@gmail.com](mailto:saboaba@gmail.com)

## INTRODUCTION

Essential oils or volatile oils are a complex mixture of compounds synthesized by living organisms and isolated from different parts of plants. Essential oils differ from plant to plant; though have a number of physical properties in common. A vast majority are known to have medicinal properties (1).

*Cedrela odorata* L. (Meliaceae) is commonly referred to as Spanish cedar wood, cigar-box tree or Brazilian mahogany. *C. odorata* is a monoecious, deciduous, and medium-sized to large tree up to 40 m tall that arrays from Mexico to South America and the West Indies (2). The tree is known for its red, rot-resistant wood that is used to make furniture, guitars (3) and cigar boxes. The repelling smell of the wood to insects makes it suitable for the making clothing chests and wardrobes. The wood from this tree is among the most sought-after in Latin America and elsewhere as it is resistant to attacks by fungi and insects, and it keeps a pleasant fragrance for many years (4).

The plant is used in traditional medicine; the decoctions of the bark are used as a remedy for wounds, fever, bronchitis, indigestion and other gastrointestinal ailments, vomiting, haemorrhage (5). *C. odorata* essential oil has been shown to have anti-microbial activity (2) and the stem extract has been found to possess anti-oxidant activity (6). Kipassa et al., 2008 (7) isolated limonoids from the stem bark. The odour from the essential from *C. odorata* is in resemblance to the odour of cedar-wood and some species of *Cyperus* oils. It is used mainly for the perfumery of soaps, some types of sprays and disinfectant (8).

The chemical compositions of the essential oils of some parts of *C. odorata* have been investigated and generally, have sesquiterpenoids in abundance (2, 5, 6, 9, and 10). However, there are no comprehensive reports on the essential from the wood of *C. odorata*.

*Dalbergia latifolia* Roxb. commonly referred to as Indian rosewood or Bombay black belongs to the Fabaceae family. The plant is usually a single stemmed tree with a dome shaped crown of lush green foliage with characteristic smell (11). It is widely distributed in tropical and subtropical regions of Bihar, Bundelkhand and Central India (12). Many *Dalbergia* species are important timber trees which are valued for their decorative and fragrant wood, rich in aromatic oils (13). Traditionally, *D. latifolia* is reported to be used as stimulant, appetizer, anthelmintic, spasmogenic, and also in the cure of dyspepsia, diarrhea, cutaneous affections and leprosy (14). It is also regarded as brain tonic to the nervous system (15). Some parts of the tree are reported to have antioxidant scavenging potential and antibacterial activities (16). Ethanol extracts of *D. latifolia* root neuropharmacological activity as nootropic and also having anxiolytic property, demonstrated in albino rats (17).

Though different phytochemical and pharmacological studies have been done on *D. latifolia*, there are no previous references in literature on the essential oils from parts of the plant.

This paper therefore, presents the comprehensive characterization of the essential oils from the leaves and woods of *C. odorata* as well as *D. latifolia* from South west Nigeria with a view to comparing any existing literature on the four oils.

## EXPERIMENTAL

Fresh samples were cut in March, 2014 from mature trees growing at the Botanical garden, University of Ibadan. They were authenticated at the Department of Botany, University of Ibadan by Mr. Donatus and the voucher specimen was deposited at the Forest Research Institute of Nigeria (FRIN) Herbarium with Voucher Number, 110039 and 110038 for *Cedrela odorata* and *Dalbergia latifolia* respectively. The stem-bark was scrapped off the wood; the fresh wood and leaves were chopped into pieces, air-dried and crushed using a fast rotating grinding machine.

### Essential oil isolation

The ground samples were subjected to the hydro-distillation method using an all glass Clevenger apparatus according to the *British pharmacopoeia* (18) method for 4 h. The oils were collected in hexane and kept in the refrigerator at 4 °C until analyses.

### Gas chromatography/ Mass spectrometry Analysis

The composition of the essential oils were determined by a Gas Chromatography- Mass spectrometry (GC-MS) using a Agilent 7890N GC Agilent mass detector Triple Quad 7000A in EI mode at 70 eV (m/z range 40 – 600 amu) and an Agilent ChemStation data system. The GC column was equipped with an HP-5MS column (30 m × 250 µm × 0.25 µm) a split-split less injector heated at 200 °C and a flame ionization detector (FID) at 230 °C. The oven temperature was programmed as follows; initial temperature 40 °C for 5 mins increased 5 °C/min to 180 °C for 6 mins and then 10 °C/min to 280 °C for 12 mins. Helium was the carrier gas used at the flow rate of 1 mL/min. The injection volume was 2 µL (split ratio 1:20).

The components were identified by comparison of their mass spectra with NIST 1998 library data of the GC-MS system as well as by comparison of their retention indices (RI) with relevant literature data (19). The relative amount of each individual component of the essential oil was expressed as the percentage of the peak area relative to the total peak area. RI values of each component were determined relative to the retention times of a homologous n-alkane series with linear interpolation on the HP-5MS column.

## RESULTS AND DISCUSSION

The chemical composition of the volatile oils of *C. odorata* leaf, *C. odorata* wood, *D. latifolia* leaf and *D. latifolia* wood are presented in **table 1**. The percentage yields of the oils (w/w) based on dry matter are 0.82%, 0.5%, 0.43% and 0.39% respectively. The four oils had different shades of yellow.

*C. odorata* leaf oil contained 10 compounds representing 97.63% of the entire oil with predominantly 85.21% sesquiterpenes with 51.65% oxygenated sesquiterpenes and 34.56% sesquiterpene hydrocarbon. The sesquiterpene hydrocarbons present are β-elemene (23.03%), α-bergamotene (4.09%), δ-cadinene (4.13%), α-humulene (2.28%) while the oxygenated sesquiterpenes are (-)-spathulenol (42.49%), cubenol (8.71%) and germacrene-D-4-ol (0.45%). Monoterpene hydrocarbon in the oil represents 8.97% as characterized by α-santalene (4.57%) and α-copaene (4.4%). The results from this study when compared

with Asekun and Ekundayo, 1999 had some variations. Asekun and Ekundayo (10) characterized 26 compounds which amount to 70.6% of the oil identified with about 30% of the constituents unidentified. Their analyses also had sesquiterpenes as the abundant compound with  $\alpha$ -santalene (9.5%), the dominant compound. Similar compounds to both studies are  $\alpha$ -copaene,  $\beta$ -elemene,  $\alpha$ -santalene,  $\alpha$ -bergamotene,  $\alpha$ -humulene, and  $\delta$ -cadinene. Variations in qualitative and quantitative compositions observed from different studies are due to seasonal and maturity variation, geographical origin, genetic variation, growth stages and postharvest drying and storage (20).

The wood of *C.odorata* had 39 compounds amounting to 99.16% of the oil; with 54.92% sesquiterpene hydrocarbon and 37.40% oxygenated sesquiterpene. The remaining 6.84% of the characterized compounds are monoterpenes and non-terpenoid compounds. The abundant sesquiterpene hydrocarbon are  $\alpha$ -cedrene (17.56%),  $\alpha$ -curcumene (12.3%),  $\alpha$ -bergamotene (5.63%), and,  $\beta$ -farnesene (3.89%) while oxygenated sesquiterpenes found in abundance are  $\beta$ -bisabolol (10.95%), z-nerolidol (9.23%),  $\gamma$ -eudesmol (8.84%,  $\alpha$ -eudesmol (5.43%). Our study also revealed the presence of  $\alpha$ -curcumene,  $\beta$ -bisabolol, juniper camphor,  $\delta$ -cadinene as observed by Motl and Trka, 1973 (8).

It is notable that the variation in the composition of the leaf and wood essentials presumably because of different parts of the plants where the samples were procured. Common constituents to the oils are  $\alpha$ -santalene,  $\alpha$ -copaene,  $\beta$ -elemene,  $\alpha$ -bergamotene, (-)-spathulenol, and  $\alpha$ -humulene. Generally, the result of this study conveniently describes *C. odorata* as a sesquiterpene chemo type thus confirming previous qualitative and quantitative analyses (2, 7, 8, 10).

The leaf oil of *D.latifolia* contained 72 compounds representing 95.13% of the oil with sesquiterpenes hydrocarbons, oxygenated sesquiterpenes and non-terpene compounds. The sesquiterpene hydrocarbon constituted 31.10% of the oil while 12.35% are oxygenated sesquiterpenes and the remaining components non-terpene compounds. The most abundant compounds are  $\alpha$ -bergamotene (4.23%,  $\alpha$ -selinene (4.89%),  $\beta$ -caryophyllene oxide (5.04%, (-)-spathulenol (4.49%), heptacosane (6.31%, and 1-heptariacotanol (5.7%).

On the other hand, 6 non-terpene compounds were identified in the wood oil of *D. latifolia* representing 95.13% of the entire oil. They are toluene (6.43%), methylcyclohexane (4.93%), ethylbenzene (11.90%), p-xylene (58.80%), m-xylene (16.71%) and allyltetramethoxybenzene (1.06%).

Three of the four essential oils; *C. odorata* leaf, *C. odorata* wood and *D. latifolia* leaf had in common (-)-spathulenol,  $\alpha$ -bergamotene and  $\beta$ -elemene. These compounds have been found to possess different biological activities (21, 22, 23, 24, 25) and is a lead to explore the bioassay of the oils especially *D. latifolia* whose essential oils have never been the subject of literature discussion.

## CONCLUSIONS

This paper provides a comprehensive characterization of the volatile oils from four different oils viz; *C. odorata* (leaf), *C. odorata* (wood), *D. latifolia* (leaf) and *D. latifolia* (wood). It thus gives a vivid comparison of the leaf oil of *C. odorata* with previously presented studies in literature and has also reported for the first time the characterization of the essential oils of *C. odorata* wood, leaf and wood oils of *D. latifolia*. The identified compounds which have been reported in some literatures to possess different biological activities lend support for the use of the two studied plants in ethno medical practice for various purposes. It is therefore worthwhile to explore the various biological properties of the essential oils with a view to ascertain their bioactivity.

**Table 1:** Chemical Constituents of The Essential Oils of *Cedrela odorata* and *Dalbergia latifolia* Leaves and Wood.

RI	CONSTITUENTS	COL	COW	DLL	DLW
677	1-Methylpyrrole	-	-	3.22	-
752	3-Methylheptane	-	-	0.09	-
760	Ethylcyclopentane	-	-	0.07	-
781	Methylcyclohexane	0.8	-	2.53	-
794	Toluene	-	-	1.26	6.43
806	Hexanal	-	-	0.12	-
819	2, 4-dimethyl-1-heptane	-	-	0.15	-
842	(z)-1, 3-dimethylcyclohexane	-	-	0.08	-
852	4-methyloctane	-	-	0.13	-
877	Methylcyclohexane	-	-	-	4.93
880	Ethycyclohexane	-	-	0.06	-
887	3,5-dimethyloctane	-	-	0.12	-
893	Ethylbenzene	-	-	1.38	11.9
905	Heptenal	-	-	0.12	-
907	p-Xylene	-	0.22	-	58.8
907	m-Xylene	-	-	-	16.71
969	1-Octen-3-ol	-	-	0.31	-
974	Vinylhexanoate	-	-	0.14	-
986	2,6-dimethylnonane	-	-	0.08	-
1040	2-pentylfuran	-	-	0.51	-
1048	(z)-2-(2-pentenyl)-furan	-	-	0.06	-
1086	2,2,6-trimethylcyclohexanone	-	-	0.17	-
1104	Nonanal	-	-	0.88	-
1112	E-2-Nonenal	-	-	0.36	-
1120	(E,Z)-2,6-Nonadienal	-	-	0.26	-
1179	(+)-dihydrocarvone	-	-	0.07	-
1186	Safranal	-	-	0.06	-
1202	Decanal	-	-	0.12	-
1204	$\beta$ -Cyclocitral	-	-	0.33	-
1211	$\alpha$ -Santalene	4.57	1.28	1.95	-
1221	$\alpha$ -Copaene	4.40	0.74	1.55	-
1231	Naphthalene	-	-	0.07	-
1281	Isoaromadendrene	-	0.40	-	-
1281	Isoaromadendrene epoxide	-	-	2.21	-
1286	Diepicedrene-1-oxide	-	-	0.90	-
1293	Aristolene epoxide	1.75	-	-	-
1303	2,6,6-trimethyl-1-Cyclohexene -1-acetaldehyde	-	-	0.17	-
1357	2-(2,6,6-trimethyl-1-cyclohexen-1-yl) ethanol	-	-	0.66	-
1377	$\delta$ -Elemene	-	-	2.38	-

# SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY

SOUTH. BRAZ. J. CHEM., Vol. 23, No. 23, 2015

88

*Ugah, T. C, Aboaba, S. A and Ekundayo, O*

1395	2-butyl-1-octanol	-	-	0.15	-
1398	$\beta$ -Elemene	23.06	2.08	3.96	-
1400	$\alpha$ -Longipinene	-	0.67	-	-
1403	$\alpha$ -Patchoulene	-	0.95	-	-
1406	(-)-aristolene	-	-	0.20	-
1410	$\alpha$ -Cedrene	-	17.56	-	-
1419	Isolatedene	-	-	0.29	-
1424	E-1,10-dimethyl-E-9-decalinol	-	-	1.31	-
1425	Epi- $\beta$ -Santalene	-	0.33	0.35	-
1429	E- $\alpha$ -Ionone	-	-	0.27	-
1430	$\alpha$ -Bergamotene	4.09	5.63	4.23	-
1431	Elocene	-	1.14	-	-
1440	$\beta$ -Farnesene	-	3.89	-	-
1446	$\beta$ -Sesquiphellandrene	-	2.43	-	-
1461	$\gamma$ -Gurjunene	-	0.90	2.09	-
1462	Aromadendrene oxide-(1)	-	-	0.68	-
1469	$\delta$ -Cadinene	4.13	-	2.80	-
1470	$\beta$ -Eudesmene	-	0.45	3.26	-
1474	$\alpha$ -Selinene	-	-	4.89	-
1480	$\delta$ -Guaiene	-	0.54	-	-
1489	Guaia-1(10),11-diene	-	-	0.72	-
1492	Isotridecylalcohol	-	-	0.16	-
1502	2-isopropenyl-4a,8-dimethyl-1,2,3 4,4a,5,6,7-octahydronaphthalene	-	-	1.2	-
1507	$\beta$ -Caryophyllene oxide	-	0.71	5.04	-
1512	Pentadecane	-	-	0.31	-
1522	Elemol	-	1.8	0.28	-
1524	$\alpha$ -Curcumene	-	12.31	1.40	-
1530	Ledol	-	0.41	0.84	-
1531	$\alpha$ -Bisabolene epoxide	-	-	0.14	-
1536	(-)-Spathulenol	42.49	1.12	4.49	-
1543	$\alpha$ -Cedrol	-	1.54	-	-
1547	$\alpha$ -Calacorene	-	-	0.2	-
1559	(E)-Longipinocarveol	-	0.52	-	-
1564	Z-Nerolidol	-	9.23	-	-
1574	Longiverbenone	-	0.22	-	-
1571	$\alpha$ -Caryophyllene	-	-	0.94	-
1579	$\alpha$ -Humulene	2.28	0.24	-	-
1580	Cubenol	8.71	-	-	-
1593	$\beta$ -Eudesmol	-	-	1.59	-
1598	$\alpha$ -Eudesmol	-	5.43	-	-
1619	$\beta$ -Bisabolol	-	10.95	-	-
1625	$\alpha$ -Bisabolol	-	2.73	-	-
1626	$\gamma$ -Eudesmol	-	8.84	-	-
1645	6-isopentyl-4,8a-dimethyl-4a,5,6,7,8,8a- hexahydro-1H-naphthalene-2-one	-	1.32	-	-
1647	Juniper Camphor	-	1.12	-	-
1660	Germacrene-D-4-ol	-	0.45	-	-

1680	Eudesm-7(11)-en-4-ol	-	-	3.87	-
1729	Murolan-3,9(11)-diene-10-peroxy	-	-	0.65	-
1739	Allyltetramethoxybenzene	-	-	-	1.06
1754	Hexahydrofarnesylacetone	-	-	2.19	-
1909	Labda-8(20),12,14-triene	-	0.50	-	-
1968	Hexadecanoic acid	-	0.25	-	-
1997	(9Z)-9,17-Octadecanal	-	0.26	-	-
2045	Phytol	-	-	0.65	-
2072	Cembrene	-	-	0.61	-
2075	Heptacosane	-	-	0.31	-
2228	(Z)-9-Octadecenamide	-	-	0.47	-
2271	Methyldehydroabietate	-	-	0.19	-
2375	Eicosylacetate	-	-	0.75	-
2704	Diisooctylphthalate	-	-	0.6	-
2705	Heptacosane	-	-	6.31	-
2788	p-Cresol,2,2-methylenebis[6-tertbutyl]-	-	5.30	-	-
2804	Octacosane	-	-	3.16	-
2944	Andrographolide	-	1.35	-	-
3942	1-Heptariacotanol	-	-	5.71	-
<b>Total</b>		<b>97.63</b>	<b>99.16</b>	<b>95.13</b>	<b>99.83</b>

COL - *Cedrela odorata* Leaves;  
DLW - *Dalbergia latifolia* wood.

COW - *Cedrela odorata* wood;  
RI - Retention Indices from present study.

DLL - *Dalbergia latifolia* leaves;

## ACKNOWLEDGEMENT

The authors are grateful for the TWAS-UNESCO Associateship awarded to Dr Sherifat A. Aboaba tenable at the H.E.J Research Institute of Chemistry, International Centre for Chemical and Biological Studies, University of Pakistan, Pakistan which provided funding for this research.

## REFERENCES

1. F. Chlodwig and N. Johannes., Sources of essential oils. *Handbook of essential oils* (2010), Vol 3: pp. 39-81.
2. E. V. Heather, A. T. Jessika, A. H. William and N. S. William, *Der Pharma Chemica*, 1(2): 14-18 (2009).
3. W. A. Zuchowski., *Guide to Tropical Plants of Costa Rica*, Zona Tropical Publishing, Miami, Florida (2005).
4. N. R. Aníbal., Part II—Species Descriptions; *Cedrela odorata* L. Instituto de Ecología, A.C. Xalapa, Veracruz, México (1986), pp. 386 – 389.
5. J. F. Morton., *Atlas of Medicinal Plants of Middle America*, vol. 1, Charles C. Thomas Publisher, Springfield, Illinois (1981).
6. R. Khaled, *Hygeia. J. D. Med.* 6(1): 25 -30 (2014).

7. N. T. Kipassa, T. Iwagawa, H. Okamura, M. Doe, Y. Morimoto and M. Nakatani., *Phytochemistry*, 69: 1782-1787 (2008).
8. O. Motl and A. Trka, *J. Sco. Cosmet. Chem.* 24, 747 (1973).
9. B. H. L. N. S. Maia, J. R. De Paula, J. Sant'ana, M. F. D. G. F. Da Silva, J. B. Fernandes, P. C. Vieira, M. D. S. S. Costa, O. S. Ohashi, J. N. M. Silva, *J. Braz. Chem. Soc.*, 11: 629-639 (2000).
10. O. T. Asekun and O. Ekundayo., *Flavour Fragr. J.* 14(6): 390-392 (1999).
11. A. G. D. Prasad, K. S. J. Cahndra and A. N. Y. Reddy., Initiation of a genetic improvement program for *Dalbergia latifolia* in Karnataka, India. Paper presented at the International *Dalbergia* workshop (1993).
12. K. R. Kirtikar and B. D. Basu., *Indian medicinal plants*. International Book Distributor, Dehradun, India (2005).
13. R. N. Chopra, S. L. Nyer and I. C. Chopra, *Supplement to the glossary of Indian medicinal plants*, CSIR, New Delhi. (1980), pp: 90.
14. M. Raji, N. Veeraiyan, T. Binu and M. Chitrabalam., *European J. environ. ecol.*, 1(1):33-43 (2014).
15. C. K. Madhava, K. Sivaji and K. Thulasi., *Flowering plants of Chittor district*, Andhra Pradesh, India, 2nd edition, (2008), Student off set printer, Tirupati.
16. D.P. Prasad, A. Pranavi, S. Kavamini and V. Suba., *JGTPS*, 4(3): 1039- 1043 (2013).
17. D. P. Prasad, C. V. Ravi C, S. Kavimani and V. Suba, *JGTPS*, 4(3): 1193-1199 (2013).
18. European Pharmacopiea 6<sup>th</sup> edition. EDQM (Council of Europe), Strasbourg, France (2008).
19. R. P. Adams., *Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy*, 4<sup>th</sup> Ed., Allured Publishing Corp., Carol Stream, IL, USA (2007)
20. S. Aboaba., Characterization and Biological Activities of Essential Oils: In: *Recent Progress in Medicinal Plant, Volume 36: Essential Oils I*. Bhattacharya, S (Eds) Studium Press, LLC, U.S.A (2013), pp. 1-20.
21. C. Salis, J. Becerra, C. Flores, J. Robledo and M. Silva., *J. Chil. Chem. Soc.*, 49(2): 157-161 (2004).
22. L. M. Leandro, F. S. Vargas, P. C. S. Barbosa and J. K. O. Neves., *Molecules*, 17: 3866-3889 (2012).
23. H. Eggenesperger: *Multiaktive Wirkstoffe fur Kosmetika Verlag fur chemische Industrie*, H. Ziolkowsky GmbH, Augsburg, (1995), pp. 34-59.
24. E. Szoke, E. Maday, J. Gershenzon, J. L. Allen and E. Lemberkovics, *J. Chromatogr. Sci.* 42: 229-234 (2004).
25. N. Boutaghane, A. Kabouche, R. Touzani, Y. A. Maklad, A. El-Azzouny, C. Bruneau and Z. Kabouche, *Nat. Prod. Comm.* 6(2): 251-252 (2011).

**CHEMICAL COMPOSITIONS AND ANTIMICROBIAL ACTIVITY OF THE LEAF  
ESSENTIAL OIL OF GOSSYPIUM HIRSUTUM**

**Ogunmola Oluranti O.<sup>1</sup>**

**<sup>1</sup>Department of Chemistry, Emmanuel Alayande College of Education, Oyo,  
Oyo State, Nigeria.**

**ABSTRACT**

*The study was conducted to analyze the chemical constituent and to evaluate the antimicrobial properties of the leaf essential oil of Gossypium hirsutum. The oil was obtained by hydrodistillation and was analyzed by gas chromatography and gas chromatography –mass spectrometry (GC/GC-MS). This revealed the presence of fifty components accounting for - % of the total oil fraction. The leaf oil was dominated by patchoulane (14.70%), Ally 1-2, 6,6 – trimethyl bicyclo (3.1.1) heptane (5.95%). 1,7, 7 – trimethyl bicyclo (2.2.1) heptanes (5.95%) and 9- (1 – methylethylidene) bicyclo (6.1.0) nonane (5.95%). The oil displayed high antimicrobial potentials to some tested micro organisms.*

**KEYWORDS:** *Gossypium hirsutum, patchoulane, aliphatic alcohol, antimicrobial potential.*

**RESUMO**

*O estudo foi conduzido para analisar o constituinte químico e avaliar as propriedades antimicrobianas do óleo essencial de folha de Gossypium hirsutum. O óleo foi obtido por hidrodestilação e analisado por cromatografia gasosa e cromatografia gasosa-espectrometria de massa (GC / GC-MS). Isso revelou a presença de cinquenta componentes representando -% da fração de óleo total. O óleo de folha foi dominado por patchoulane (14,70%), Ally 1-2, 6,6 - trimetilbicyclo (3.1.1) heptano (5,95%). 1,7, 7-trimetilbicyclo (2.2.1) heptanos (5,95%) e 9- (1-metiletilideno) bicyclo (6.1.0) nonano (5,95%). O óleo apresentou potenciais antimicrobianos elevados para alguns microrganismos testados.*

**PALAVRAS CHAVE:** *Gossypium hirsutum, patchoulene, álcool alifático, potencial antimicrobiano*

**Introduction**

Plants are one of the five main groups of living things, these groups are called kingdoms. The plant kingdom includes such living things as grasses, trees, ferns, bushes and flowers. About 260,000 species of plant exist on the earth (14) plant contains complex and powerful substances known as essential oils. These are aromatic liquid derived from shrubs, flowers, trees, roots, bushes, herbs and seeds (15). Medicinal plants form an aspect of cultural heritage of the human race and their use is still expanding. Plants with useful medicinal properties have been obtained from folklore records of many indigenous communities. Egyptian and Chinese priests and physicians since time immemorial had been using extracted oils from plant for healing and this makes essential oil the earliest known medicine (15). An essential oil is a concentrated hydrophobic liquid containing volatile aroma compounds from plants. It is also known as volatile oils, ethereal oils or aethrolea or simply as the "oil of" the plant from which they were extracted. An oil is "essential" in the sense that it carries a distinctive scent or essence of the plant (16).

*Gossypium hirsutum* L. (family malvaceae) is a cotton specie native to Central America and the Caribbean. After being domesticated in the United States, this cotton species now provides over 90% of commercial cotton worldwide (13). In Nigeria, cotton seed provides raw materials for local spinning and weaving industry (4). It is believed to be an indigenous effective medicinal plant. The dried root bark of cotton contains gossypol, this compound may cause abortion – inducing effect (3). Gossypol is a polyphenol that may be isolated more easily from the bark (5). There is on going research to use gossypol as an alternative to vasectomy (7), the plant is known to

posses anti-malaria properties and it has demonstrated in vitro effects on cell regulation and antitumor activity against mammary carcinoma a cell lines (9, 18, 6).

When the chemical constitution of the essential oil of six species within three genera of the family malvaceae were investigated, the essential oil of *Gossypium tursutum* L. was found to contain 67.0% hydrocarbon and beta-bisabolol was the major oxygenated component 13.7% (1), also (12) reported the presence of fifty nine constituents in the oil aroma of *Gossypium barbadense* in which patchoulane (7.94%), Bicyclo (4.1.0) heptanes -3-cylopropy 1-7-hydroxymethyl (trans) (7.86%), Z-Nonadecatetraene (6.19%), alpha-patchoulane (6.19%), caryophyllene oxide (5.75%) and Bicyclo (7.2.0) undec-4-ene, 11, 11-trimethy 1-8-methylene (IR\*, 4z, 95\*) (3.50%) as the major constituents. The oil also has antibiotic potential (12, 13).

## **Materials and Methods**

### **Plant Material**

Leaves of *Gossypium hirsutum* were collected from Oyo, south West, Nigeria. The plant was authenticated at the Forest Research Institute of Nigeria (FRIN) Ibadan, Nigeria. Extraction of the *hirsutum* was air dried in a well ventilated place till when the moisture content reduced to a minimum suitable for grinding. The plant material was pulverized and used immediately. The crushed plant material (400g) was subjected to hydrodistillation for 3 hours using a Clevenger – type apparatus (11). The oil collected was stored in a vial at low temperature until the time of the analysis.

### **Multidimensional GC X GC-MS Analysis**

Analysis of leaf essential oil of *Gossypium barbadense* was performed using multidimensional chromatogram coupled with gas chromatography, mass spectrometer (Shimadzu, Japan,) equipped with double capillary columns (2.50m X 0.25<sup>N</sup>m I.d, 0.25<sup>N</sup>mdf) that have different characteristics (non polar and polar) High purity helium was used as the carrier gas at a constant flow rate of 0.99ml/min. A total of 1 $\mu$ l sample was injected (Split ratio 100:1) into GC X GC MS using AOC 20i auto injector for analysis. The initial temperature was set at 60<sup>o</sup>C treated at a rate of 30<sup>o</sup>C minute to 28<sup>o</sup>C and held isothermally for 6 minutes. Ion source temperature for this analysis was set at 200<sup>o</sup>C while the interface temperature was set at 250<sup>o</sup>C solvent cut time was 3.0 minutes and the mass spectrometer was set to operate in electron ionization mode with an ionizing energy of 70eV as acquisition mass range from 40 – 700 a.m.u at 0.50cscan/S.

### **Antimicrobial Activities**

The antibacterial activity of the leaf essential oil of *Gossypium barbadense* was measured against Gram-negative (*Escherichia coli*, *Klebsiella pneumoniae*, *Salmonella typhimourion* and *Proteus mirabilis*) and Gram – positive (*Staphylococcus aureus* and *Streptococcus agalactiae*) using a well diffusion method according to the National Committee for Clinical Laboratory Standard (13). Petri plates containing approximately 25-30ml of nutrient agar medium were swabbed using cotton applicator with 24 hours sub-cultured bacteria strains which were prepared in dilution to match the turbidity intensity of the MacFarland standard. Wells (6mm diameter) were punched in the agar

and filled 10<sup>N</sup>L of the extract of different concentration (1000, 100 and 10Ngml<sup>-1</sup>). The plates were incubated at 37<sup>0</sup>C for 24 hours. The antibacterial activities were assessed by measuring the inhibition zone diameter (mm) around the well.

### **Results and Discussion**

Exhaustive hydro distillation of the leaf of *Gossypium hirsutum* gave a light yellow coloured oil with 0.4% V/W yield per 400g of dried leaf sample. The analysis of the leaf essential oil was carried out using GC X GC – MS and the percentage compositions are given in table 1. Fifty compounds were identified from the leaf essential oil amounting to –% of the oil. The leaf essential oil was dominated by patchoulane (14.70%) Allyl – 2, 6, 6 – trimethyl bicyclo (3.1.1) heptanes (5.95%), 1, 7, 7- Trimethyl bicyclo (2.2.1) heptanes (5.95%) and 9-(1-methyl-ethylidene) bicyclo (6.1.0) nonane (5.95%). The composition of the essential oil has similarity with the constituent of *Gossypium barbadense* reported by (12), patchovlane, been the compound with the highest percentage in *Gossypium barbadense* has the highest percentage in *Gossypium hirsutum*(14.70%). The aliphatic reported by (12) as part of the constituents of *Gossypium barbadense* are also in the present study even though it is quantitatively different (6 – Mehthyl-2-Vingl-5-hepten-01 (1.30%), 2- pentadecen-1-01 (1.30%) and Z – 10-pentadecen-1-01.

In addition, the oil displayed high antimicrobial activities to proteus micrabelis at 1,000, 100 and 10 Ngml<sup>-1</sup>, Esherichia coli at 1,000, 100 and 10Ngml<sup>-1</sup>, staphylococcus aureus at 1,000, 100 and 10Ngml<sup>-1</sup> and klebsiell pneumonia at 1,000, 100 and 10Ngml<sup>-1</sup> as shown in table 2. The oil also displayed low inhibitory activity against gram negative

salmonella typhimurium at all concentrations and this can antibiotics as shown in table

3.

### Conclusion

The result obtained in this study showed that *Gossypium hirsutum* possess essential oil and it has a great antibiotics potential and this confirm its use in traditional medicine as an antibiotics and anti-malaria.

**Table I**

Chemical Constituents of the essential oil of *Gossypium Hirsutum*.

S/N	R.I	Compound	%Comp
1.	1061	2 (3H) – Furanone	2.00
2.	837	2 – Pentanethol	4.00
3.	935	Nitrohexane	2.00
4.	919	Pyruvic acid, Methylester oxime	2.00
5.	620	3 – Ethoxy-1-butene	0.58
6.	1273	3, 5 – Octanedione, 2, 2, 4, 7 – tetramethyl	0.58
7.	821	Pentanal, 2, 2 – dimethyl	0.58
8.	676	Tert – butylcarbinol	0.58
9.	729	Methyl -2-2methoxyacrylate	0.58
10.	1515	Germacrene	0.2
11.	1221	Copane	0.4
12.	1440	Alpha – Amorphene	0.2

13.	1458	(Z, Z) alpha – Farnesene	0.2
14.		2, 6 – Bis (diazn) adamantane	0.32
15.	1050	2, 8 – Decadiyne	0.32
16.	1229	Ethanone-1-2-5 (methyl)-2 forylcycloprophyl	0.32
17.	2139	Benzene, 2-phenyl-2-propenyl) sulfoni oxacyclo tetradeca-4, 11-ddiyne	0.32
18.	1393	Patcheulane	14.70
19.	1148	1-Methyl-2-methylene-3, 5-divinyl cyclohexane	1.40
20.	1494	4,11, 11-Trimethyl-8-methylene bicyclohexane	1.40
21	1369	Cyclohexene, 4-180 propenyl-1-methoxymethyl	1.40
22.	1579	Alpha Humulene	0.63
23.	958	Ocimene	1.26
24.	976	Beta-CIS-Ocimene	1.26
25.		3-Methyl-4-(phenylthia)-2-prop-2-enyl-2, phenel, 1-dioxide	0.57
26.		Lauryl pyridinium chloride	0.57
27.	1128	Pentanol-5- (methylene Cyclopropyl)	0.57
28.	1934	Z, Z, Z – 4, 6, 9 – Nonadecatriene	0.57
29.	1414	1, 2-Epoxy – 5, 9 – Cyclododecadiene	0.57
30	1187	Allyl – 2, 6, 6 – trimethyl bicoclo (3.1.1) heptanes	5.95
31.	1326	1, 7, 7 – trimethyl bicyclo(2.2.1) heptanes	5.95

# SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY

SOUTH. BRAZ. J. CHEM., Vol. 23, No. 23, 2015

98

Ogunmola Oluranti O.

32.	1242	9 - (1 – Methylethylidene) bicycle (6.1.0) nonane	5.95
33.	1066	1-(1-Methylene-2-propenyl) cyclopentanol	1.56
34.	1307	Bicyclo (4.1.0) heptanes-3-cyclopropyl 1-7-hydroxymethyl trans	1.56
35.	1307	Bicyclo (4.1.0) heptanes-3-cyclopropyl 1-7-hydroxymethyl cis	1.56
36.	1078	1-(2 –Methylene cyclopropyl) cyclopentanol	1.56
37.	1143	Ethylidene cyclo octane	1.56
38.	1196	Dihydrocarveol	1.30
39.	1169	6 – Methyl – 2-Vinyl-5-hepten-1-01	1.30
40.	1772	2 – Pentadecyn-1-01	1.30
41.	1292	Myrcenylacetate	1.30
42.	1079	Isotujol	1.30
43.	1968	Palmitic acid	3.27
44.	1272	Pelargic acid	1.09
45.	1769	Myristic acid	1.09
46.	2175	Cis-Oleic acid	4.00
47.	1763	Z-10-pentadecen-1-01	4.00
48.	2007	9 – Octadecenal	4.00
49.	1763	E-10-Pentadecen-1-01	4.00
50.	1293	10 – Undecenal	4.00
		Total	

**Table 2**

Zones of inhibition (mm) showing the antimicrobial activities of gram-negative and gram-positive bacteria.

Organism	Escherichia Coil	Klebsiella Pneumonia	Salmonella Typhimouriu m	Proteus Mirabelis	Staphylococos	streptococcus
Conc. (Ngml <sup>-1</sup> )	1,000 100 10	1,000	1,000 100 10	1,000 100 10	1,000 100 10	1,000 100 10
Zone of inhibition	20 17 10	15 15 15	09 08 -	32 30 24	20 20 20	11 11 11

**Keynote:** - = no inhibition, 6-9mm = low inhibition, 10 – 15mm = moderate inhibition

And ≥ 15mm = high inhibition

+ = gram-negative bacteria

\* = gram-positive bacteria

**Table 3**

Zones of inhibition (mm) showing the antimicrobial activities of antibiotics against gram positive and gram negative bacteria.

Organism/ antibiotics/ concentration	Escherichia Coil	Klebsiella Pneumonia	Salmonella Typhimouriu m	Proteus Mirabelis	Staphylococcus aureus	streptococcus agalactiae
Nitroforantion (2Ng)	25	20	15	15	28	11
Ofloxacin (5Ng)	20	20	22	33	20	-
Gentamicin (10Ng)	24	15	18	16	-	-

**Keynote:** - Same as in table 2.

## References

1. Thompson, A. C., Hanny, B. W., Hedin, P. A. and Gueldner, R. C. (1971) Phytochemical Studies in the family malvaceae 1. Comparison of essential oils of six species by gas – liquid chromatography. *American Journal of Botany* 58: 9. Pp 803-907.
2. Gossypium Tincture from Amazon herbs. Retrieved from Tropilan inc., exporter & wholesaler of medicinal plants herbs and tropical seeds on 27-05-15.
3. [www.naturalstandard.com](http://www.naturalstandard.com). Accessed 27<sup>th</sup> May 2015.
4. Esssien, E. E., Aboaba, S. O. and Ogunwande, I. A. (2011) Constituents and Antimicrobial properties of the leaf essential oil of *Gossypium barbadense* (linn). *Journal of Medicinal Plant Research* 5 (5) Pp 702-705.
5. Zhou, R. H., and Lin, X. D. (1988) Isolation of (-) –gossypol from natural plant. *Contraceptives*, 37: Pp 239 – 245.
6. Ajaiyeoba, E. O., Falade, C. O., Fawole, O. I., Akinboye, D. O., Gbotosho, G. O., Bolaji, O. M., Ashidi, J. S., Abiodun, O. O., Osowole, O. S., Itiola, O. A., Oladepo, O., Sowumi, A. and Oduola, A. M, (2004) Efficacy of herbal remedies used by herbalist in Oyo State Nigeria for treatment of Plasmodium falciparum Infections – a survey and observation. *African Journal of Medicinal Science* 33 (2) Pp 115-199.
7. Qian, S. Z. and Wang, Z.G. (1984) Gossypol: a potential antifertility agent for males. *Animal Review of Pharmacological Toxicology* 24: 329 – 360.
8. Liu, G. Z, Lyle, K. C and CaOJ. (1987) Clinical trail of gossypol as a male contraceptive drug. Part I. efficacy study. *Fertile. Steril* 48 93) 459 -461.
9. Shidaifat, F. (1996) Inhibition of human postrate cancer cells growth by gossypol Associated with stimulation of transforming growth factor-beta. *Cancer letter* 107 (1): 37-44.
10. National Committee for Clinical Laboratory standards (1993) Retrieved online from [www.nchi.nlm.gov/pmc.on](http://www.nchi.nlm.gov/pmc.on) 17-07-14.
11. European Pharmacopoeia Commission (2004). *European Pharmacopoeia* 5<sup>th</sup> Ed. Council of Europe: Strasbourg Cedex, France.
12. Ogunmola, O. O. and Larayetan, R. A. (2015) Chemical and medicinal potentials of the leaf essential oil of *Gossypium barbadense* L. *Brazilian Journal of Chemistry*.
13. Chan, B. G, Waiss Jr. A. C, Lukefahr, M. J (1978) Condensed tuning an antibiotic Chemical from *Gossypium hirsutum*. *Journal of insect physiology* 24: 113 – 118.
14. Microsoft Encarta, (2009)
15. Smyrna (2006)
16. Schnaublet (1999)
17. [www.naturalstandard.com](http://www.naturalstandard.com)
18. Salako, O. A. and Awodele, O. (2012) Evaluation of the antimalaria activity of the aqueous leaf extract of *Gossypium barbadense* (Malvaceae) in mice. *Drugs and Therapy studies* 2 (2) Pp 1-7.

BOOKS / LIVROS

**E**nciclopédia dos minerais do Brasil abrange praticamente todas as substâncias minerais existentes em nosso território e suas localizações. Dividida em sete volumes, traz as seguintes temáticas: Elementos Nativos e Halogenetos; Sulfetos e Sulfossais; Óxidos e Hidróxidos; Carbonatos, Sulfatos e Combinações Orgânicas; Fosfatos, Arsenatos e Vanadatos; Silicatos.



Próximo lançamento:



[www.editoraulbra.com.br](http://www.editoraulbra.com.br)

Paulo César Pereira das Neves, natural de São Gabriel/RS, é geólogo (1986) pela Universidade do Vale do Rio dos Sinos (Unisinos), mestre em Ciências (1992) e doutor em Geociências (1996) pela Universidade Federal do Rio Grande do Sul (UFRGS). Pós-doutor em Mineralogia (2013) pela Universidade de São Paulo (USP). É autor dos livros *Introdução à mineralogia prática* (4ª edição), *Fundamentos de cristalografia* (2ª edição), *Glossário de paleontologia e termos associados* e *Enciclopédia dos minerais do Brasil* (quatro volumes já publicados), todos pela Editora da ULBRA. É professor do curso de Química Industrial nas disciplinas de Cristalografia e Mineralogia Industrial e responsável pelo Laboratório de Geologia e Mineralogia da Universidade Luterana do Brasil (ULBRA). Também atua no curso de Pós-Graduação em Engenharia dos Materiais e Processos Sustentáveis da mesma instituição. É autor de artigos nacionais e internacionais e revisor *ad hoc* das revistas *Ciências Ambientais da Unilasalle* (Canoas), *Gaea da Unisinos* (São Leopoldo) e *Pesquisas em Geociências* da UFRGS (Porto Alegre). Também é consultor científico da Fundect (MS).



Daniel Atencio, natural de São Caetano do Sul/SP, é geólogo (1982), mestre (1986), doutor (1991) e livre-docente (1999) pela Universidade de São Paulo (USP). É professor do Instituto de Geociências da USP desde 1984, sendo o representante do Brasil na Commission on New Minerals and Mineral Names (CNMMN)/ Commission on New Minerals, Nomenclature and Classification (CNMNC) da International Mineralogical Association (IMA) desde 1990. É autor de 32 novos minerais-tipo aprovados pela IMA. O mineral atencioíta foi nomeado em sua homenagem. É autor dos livros *Type Mineralogy of Brazil* (2000) e *Enciclopédia dos minerais do Brasil* (quatro volumes já publicados), além de capítulos de livro, artigos em periódicos especializados, trabalhos em anais de eventos, resenhas, traduções, prefácios, etc. Foi editor da *Revista Brasileira de Geociências* e, atualmente, é relator de várias publicações. É supervisor do Laboratório de Difractometria de Raios X do Instituto de Geociências da USP. Tem orientado dissertações de mestrado, teses de doutorado e pós-doutorado, além de grande número de trabalhos de iniciação científica e trabalhos de conclusão de curso na área de Geociências. Atualmente, coordena projetos de pesquisa na Fapesp.



**ENCICLOPÉDIA DOS  
MINERAIS DO BRASIL**  
CARBONATOS, SULFATOS  
E COMBINAÇÕES ORGÂNICAS

**Paulo César Pereira das Neves**  
**Daniel Atencio**



Editora da ULBRA

# Sumário

<b>Prefácio .....</b>	<b>11</b>
<b>Apresentação .....</b>	<b>13</b>
<b>Disponibilização de imagens .....</b>	<b>15</b>
<b>1 Introdução .....</b>	<b>17</b>
<b>2 A classificação de Strunz e Nickel .....</b>	<b>23</b>
<b>Classe 5: Carbonatos e Nitratos .....</b>	<b>23</b>
<b>3 Carbonatos .....</b>	<b>25</b>
<b>5.A. Carbonatos sem ânions adicionais e sem H<sub>2</sub>O.....</b>	<b>25</b>
<b>5.AB. Carbonatos com cátions alcalino-terrosos e outros íons divalentes. ....</b>	<b>25</b>
3.1 CALCITA CaCO <sub>3</sub> – 5.AB.05. ....	25
3.2 MAGNESITA MgCO <sub>3</sub> – 5.AB.05.....	44
3.3 SIDERITA FeCO <sub>3</sub> – 5.AB.05. ....	48
3.4 RODOCROSITA MnCO <sub>3</sub> – 5.AB.05.....	54
3.5 SMITHSONITA ZnCO <sub>3</sub> – 5.AB.05.....	56
3.6 DOLOMITA CaMg(CO <sub>3</sub> ) <sub>2</sub> – 5.AB.10. ....	58
3.7 ANKERITA CaFe <sup>2+</sup> (CO <sub>3</sub> ) <sub>2</sub> – 5.AB.10.....	68
3.8 KUTNOHORITA CaMn <sup>2+</sup> (CO <sub>3</sub> ) <sub>2</sub> – 5.AB.10. ....	74
3.9 ARAGONITA CaCO <sub>3</sub> – 5.AB.15. ....	76
3.10 ESTRONCIANITA SrCO <sub>3</sub> – 5.AB.15. ....	79
3.11 WITHERITA BaCO <sub>3</sub> – 5.AB.15. ....	81
3.12 CERUSSITA PbCO <sub>3</sub> – 5.AB.15.....	82
3.13 HUNTITA CaMg <sub>3</sub> (CO <sub>3</sub> ) <sub>4</sub> – 5.AB.25.....	85
3.14 NORSETHITA BaMg(CO <sub>3</sub> ) <sub>2</sub> – 5.AB.25. ....	87
3.15 BARITOCALCITA BaCa(CO <sub>3</sub> ) <sub>2</sub> – 5.AB.30.....	89
3.16 CARBOCERNAÍTA (Ca,Na)(Sr,Ce,Ba)(CO <sub>3</sub> ) <sub>2</sub> – 5.AB.35. ....	90
<b>5.AC. Carbonatos alcalinos e alcalino-terras raras.....</b>	<b>92</b>
3.17 EITELITA Na <sub>2</sub> Mg(CO <sub>3</sub> ) <sub>2</sub> – 5.AC.05a.....	92
<b>5.AD. Carbonatos com elementos terras raras. ....</b>	<b>93</b>
3.18 BURBANKITA (Na,Ca) <sub>3</sub> (Sr,Ba,Ce) <sub>3</sub> (CO <sub>3</sub> ) <sub>5</sub> – 5.AD.10. ....	93

<b>5.B. Carbonatos com ânions adicionais e desprovidos de água.....</b>	<b>95</b>
<b>5.BA. Carbonatos com Cu, Co, Ni, Zn, Mg, Mn. ....</b>	<b>95</b>
3.19 AZURITA $\text{Cu}^{2+}_3(\text{CO}_3)_2(\text{OH})_2 - 5.\text{BA}.05.$ .....	95
3.20 MALAQUITA $\text{Cu}^{2+}_2(\text{CO}_3)(\text{OH})_2 - 5.\text{BA}.10.$ .....	99
3.21 HIDROZINCITA $\text{Zn}_5(\text{CO}_3)_2(\text{OH})_6 - 5.\text{BA}.15.$ .....	105
3.22 AURICALCITA $(\text{Zn},\text{Cu}^{2+})_5(\text{CO}_3)_2(\text{OH})_6 - 5.\text{BA}.15.$ .....	107
<b>5.BD. Carbonatos com elementos terras raras. Os grupos dessa subdivisão são arranjados em ordem crescente da relação entre os raios atômicos F, O, OH:CO<sub>3</sub>.....</b>	<b>109</b>
3.23 SINQUISITA-(Ce) $\text{Ca}(\text{Ce},\text{La})(\text{CO}_3)_2\text{F} - 5.\text{BD}.20.$ .....	109
3.24 SINQUISITA-(Y) $\text{Ca}(\text{Y},\text{La})(\text{CO}_3)_2\text{F} - 5.\text{BD}.20.$ .....	110
3.25 PARISITA-(Ce) $\text{Ca}(\text{Ce},\text{La})_2(\text{CO}_3)_2\text{F}_2 - 5.\text{BD}.30.$ .....	112
3.26 PARISITA-(La) $\text{Ca}(\text{La},\text{Nd},\text{Ce})_2(\text{CO}_3)_2\text{F}_2 - 5.\text{BD}.30.$ .....	114
3.27 BASTNÁSITA-(Ce) $\text{CaCO}_3\text{F} - 5.\text{BD}.35.$ .....	115
3.28 TORIOBASTNÁSITA $\text{Th}(\text{Ca},\text{Ce})(\text{CO}_3)_2\text{F}_2 \cdot 3\text{H}_2\text{O} - 5.\text{BD}.35.$ .....	118
<b>5.BE. Carbonatos com cátions metálicos: Pb e Bi. ....</b>	<b>119</b>
3.29 FOSGENITA $\text{Pb}_2(\text{CO}_3)\text{Cl}_2 - 5.\text{BE}.15.$ .....	119
3.30 BISMUTITA $\text{Bi}_2\text{O}_2\text{CO}_3 - 5.\text{BE}.20.$ .....	121
<b>5.C. Carbonatos desprovidos de ânions adicionais e com H<sub>2</sub>O.....</b>	<b>123</b>
<b>5.CA. Carbonatos com cátions de raios atômicos médios.....</b>	<b>123</b>
3.31 NESQUEHONITA $\text{MgCO}_3 \cdot 3\text{H}_2\text{O} - 5.\text{CA}.05.$ .....	123
<b>5.CB. Carbonatos com cátions de raios atômicos grandes (álcali e álcali-terras raras). ....</b>	<b>125</b>
3.32 TRONA $\text{Na}_3\text{H}(\text{CO}_3)_2 \cdot 2\text{H}_2\text{O} - 5.\text{CB}.05.$ .....	125
<b>5.CC. Carbonatos com elementos terras raras.....</b>	<b>126</b>
3.33 TENERGITA-(Y) $\text{Y}_2(\text{CO}_3)_3 \cdot 2 \cdot 3\text{H}_2\text{O} - 5.\text{CC}.10.$ .....	126
3.34 LANTANITA-(Nd) $(\text{Nd},\text{La})_2(\text{CO}_3)_3 \cdot 8\text{H}_2\text{O} - 5.\text{CC}.20.$ .....	128
3.35 LANTANITA-(La) $(\text{La},\text{Nd})_2(\text{CO}_3)_3 \cdot 8\text{H}_2\text{O} - 5.\text{CC}.20.$ .....	129
<b>5.D. Carbonatos com ânions adicionais providos de H<sub>2</sub>O.....</b>	<b>131</b>
<b>5.DA. Cátions com raios atômicos médios. ....</b>	<b>131</b>
3.36 HIDROMAGNESITA $\text{Mg}_5(\text{CO}_3)_4 \cdot 4\text{H}_2\text{O} - 5.\text{DA}.05.$ .....	131
3.37 KAMBALDAÍTA $\text{NaNi}_2(\text{CO}_3)_3 \cdot 3\text{H}_2\text{O} - 5.\text{DA}.20.$ .....	133
3.38 QUINTINITA $\text{Mg}_6\text{Al}_2(\text{OH})_{12}(\text{CO}_3) \cdot 3\text{H}_2\text{O} - 5.\text{DA}.40.$ .....	134
3.39 STICHTITA $\text{Mg}_6\text{Cr}_2(\text{OH})_{16}[\text{CO}_3] \cdot 4\text{H}_2\text{O} - 5.\text{DA}.50.$ .....	136
3.40 REEVESITA $\text{Ni}_3\text{Fe}^{2+}_2(\text{OH})_{16}[\text{CO}_3] \cdot 4\text{H}_2\text{O} - 5.\text{DA}.50.$ .....	138
<b>5.DC. Cátions com raios atômicos grandes.....</b>	<b>139</b>
3.41 CALCIOANCILITA-(Ce) $(\text{Ce},\text{Ca},\text{Sr})(\text{CO}_3)_2(\text{OH},\text{H}_2\text{O}) - 5.\text{DC}.05.$ .....	139
3.42 ANCILITA-(Ce) $\text{CeSr}(\text{CO}_3)_2(\text{OH}) \cdot \text{H}_2\text{O} - 5.\text{DC}.05.$ .....	141
<b>5.E. Carbonatos de uranila. ....</b>	<b>143</b>
<b>5.EA. Carbonatos com relação <math>\text{UO}_2:\text{CO}_3 = 1:1.</math>.....</b>	<b>143</b>
3.43 RUTHERFORDINA $\text{UO}_2\text{CO}_3 - 5.\text{EA}.05.$ .....	143

5.EE. Carbonatos com relação $UO_2:CO_3 = 1:5$ .....	145
3.44 ASTROCIANITA-(Ce) $Cu_2Ce_2(UO_2)(CO_3)_5(OH)_2 \cdot 1,5H_2O - 5.EE.05$ .....	145
<b>4 Nitratos.....</b>	<b>147</b>
Divisão 5.N: Nitratos .....	147
<b>5.NA. Nitratos sem <math>OH^-</math> e desprovidos de <math>H_2O</math>.....</b>	<b>147</b>
4.1 NITRATINA $NaNO_3 - 5.NA.05$ .....	147
4.2 NITRO $KNO_3 - 5.NA.10$ .....	149
<b>5.NC. Nitratos com <math>H_2O</math>.....</b>	<b>152</b>
4.3 NITROCALCITA $Ca(NO_3)_2 \cdot 4H_2O - 5.NC.10$ .....	152
<b>CLASSE 7 – Sulfatos, Cromatos, Molibdatos e Wolframatos.....</b>	<b>154</b>
<b>5 Sulfatos.....</b>	<b>155</b>
<b>7.A. Sulfatos, Selenatos e Teluratos desprovidos de ânions adicionais e <math>H_2O</math>.....</b>	<b>155</b>
<b>7.AC. Sulfatos com cátions de raios iônicos médios a grandes.....</b>	<b>155</b>
5.1 LANGBEINITA $K_2Mg_2(SO_4)_3 - 7.AC.10$ .....	155
5.2 THENARDITA $Na_2SO_4 - 7.AC.25$ .....	156
<b>7.AD. Sulfatos com cátions de raios atômicos grandes.....</b>	<b>158</b>
5.3 GLAUBERITA $Na_2Ca(SO_4)_3 - 7.AD.25$ .....	158
5.4 ANIDRITA $CaSO_4 - 7.AD.30$ .....	160
5.5 CELESTITA $SrSO_4 - 7.AD.35$ .....	163
5.6 BARITA $BaSO_4 - 7.AD.35$ .....	164
5.7 ANGLESITA $PbSO_4 - 7.AD.35$ .....	172
5.8 OLSACHERITA $Pb_2(SeO_4)(SO_4) - 7.AD.35$ .....	174
<b>7.B. Sulfatos (Selenatos) com ânions adicionais desprovidos de água.....</b>	<b>176</b>
<b>7.BB. Sulfatos com cátions de raios atômicos médios.....</b>	<b>176</b>
5.9 ANTLERITA $Cu^{2+}_3(SO_4)(OH)_4 - 7.BB.15$ .....	176
5.10 BROCHANTITA $Cu^{2+}_4(SO_4)(OH)_6 - 7.BB.35$ .....	177
<b>7.BC. Sulfatos com raios atômicos médios a grandes.....</b>	<b>180</b>
5.11 ALUNITA $KAl_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	180
5.12 NATROJAROSITA $NaFe^{3+}_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	183
5.13 JAROSITA $KFe^{3+}_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	184
5.14 HIDRONIOJAROSITA $(H_3O)(Fe^{3+})_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	186
5.15 PLUMBOJAROSITA $Pb_{0,5}Fe^{3+}_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	188
5.16 OSARIZAWAÍTA $Pb(Al,Cu)_3(SO_4)_2(OH)_6 - 7.BC.10$ .....	189
5.17 LINARITA $PbCu^{2+}(SO_4)(OH)_2 - 7.BC.35$ .....	190
<b>7.C. Sulfatos desprovidos de ânions adicionais e com <math>H_2O</math>.....</b>	<b>192</b>
<b>7.CB. Somente com cátions com raios atômicos médios.....</b>	<b>192</b>
5.18 KIESERITA $MgSO_4 \cdot H_2O - 7.CB.05$ .....	192
5.19 ROZENITA $Fe^{2+}SO_4 \cdot 4H_2O - 7.CB.15$ .....	193
5.20 CALCANTITA $Cu^{2+}SO_4 \cdot 5H_2O - 7.CB.20$ .....	195

5.21 MELANTERITA $\text{Fe}^{2+}\text{SO}_4 \cdot 7\text{H}_2\text{O} - 7.\text{CB}.35$ .....	196
5.22 EPSOMITA $\text{MgSO}_4 \cdot 7\text{H}_2\text{O} - 7.\text{CB}.40$ .....	199
5.23 ALUNOGÊNIO $\text{Al}_2(\text{SO}_4)_3 \cdot 17\text{H}_2\text{O} - 7.\text{CB}.45$ .....	201
5.24 COQUIMBITA $\text{Fe}^{3+}_2(\text{SO}_4)_3 \cdot 9\text{H}_2\text{O} - 7.\text{CB}.50$ .....	203
5.25 PARACOQUIMBITA $\text{Fe}^{3+}_2(\text{SO}_4)_3 \cdot 9\text{H}_2\text{O} - 7.\text{CB}.50$ .....	205
5.26 RÖMERITA $\text{Fe}^{2+}\text{Fe}^{3+}_2(\text{SO}_4)_4 \cdot 14\text{H}_2\text{O} - 7.\text{CB}.55$ .....	206
5.27 PICKERINGITA $\text{MgAl}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O} - 7.\text{CB}.60$ .....	208
5.28 HALOTRIQUITA $\text{Fe}^{2+}\text{Al}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O} - 7.\text{CB}.60$ .....	210
5.29 BILINITA $\text{Fe}^{2+}\text{Fe}^{3+}_2(\text{SO}_4)_4 \cdot 22\text{H}_2\text{O} - 7.\text{CB}.60$ .....	213
<b>7.CC. Sulfatos com cátions de raios atômicos médios a grandes.....</b>	<b>214</b>
5.30 CALINITA $\text{KAl}(\text{SO}_4)_2 \cdot 11\text{H}_2\text{O} - 7.\text{CC}.15$ .....	214
5.31 TSCHERMIGITA $(\text{NH}_4)\text{Al}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O} - 7.\text{CC}.20$ .....	216
5.32 ALÚMEN-(K) $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O} - 7.\text{CC}.20$ .....	217
5.33 LEONITA $\text{K}_2\text{Mg}(\text{SO}_4)_2 \cdot 4\text{H}_2\text{O} - 7.\text{CC}.55$ .....	219
5.34 POLIALITA $\text{K}_2\text{Ca}_2\text{Mg}(\text{SO}_4)_4 \cdot 2\text{H}_2\text{O} - 7.\text{CC}.65$ .....	220
<b>7.CD. Sulfatos com cátions de raios atômicos grandes. ....</b>	<b>222</b>
5.35 GIPSITA $\text{CaSO}_4 \cdot 2\text{H}_2\text{O} - 7.\text{CD}.30$ .....	222
<b>7.D. Sulfatos com ânions adicionais e <math>\text{H}_2\text{O}</math>. ....</b>	<b>227</b>
<b>7.DB. Sulfatos com cátions de raios atômicos médios; octaedros insulares e grupos finitos de octaedros. ....</b>	<b>227</b>
5.36 HOHMANNITA $\text{Fe}^{3+}_2\text{O}(\text{SO}_4)_2 \cdot 8\text{H}_2\text{O} - 7.\text{DB}.20$ .....	227
5.37 MAGNESIOCOPIAPITA $\text{MgFe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2 \cdot 20\text{H}_2\text{O} - 7.\text{DB}.25$ .....	228
5.38 COPIAPITA $\text{Fe}^{2+}\text{Fe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2 \cdot 20\text{H}_2\text{O} - 7.\text{DB}.25$ .....	230
5.39 ALUMINOCOPIAPITA $(\text{Al}_{2/3}\square_{1/3})\text{Fe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2 \cdot 20\text{H}_2\text{O} - 7.\text{DB}.25$ .....	232
5.40 FERRICOPIAPITA $(\text{Fe}^{3+}_{2/3}\square_{1/3})\text{Fe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2 \cdot 20\text{H}_2\text{O} - 7.\text{DB}.25$ .....	233
<b>7.DC. Sulfatos com cátions de raios atômicos médios; cadeias compatilhando as arestas dos octaedros. ....</b>	<b>235</b>
5.41 BUTLERITA $\text{Fe}^{3+}(\text{SO}_4)(\text{OH}) \cdot 2\text{H}_2\text{O} - 7.\text{DC}.05$ .....	235
5.42 FIBROFERRITA $\text{Fe}^{3+}(\text{SO}_4)(\text{OH}) \cdot 5\text{H}_2\text{O} - 7.\text{DC}.05$ .....	236
5.43 BOTRIOGÊNIO $\text{MgFe}^{3+}_2(\text{SO}_4)_2(\text{OH}) \cdot 7\text{H}_2\text{O} - 7.\text{DC}.10$ .....	238
<b>7.DD. Sulfatos com cátions de raios atômicos médios; folhas de octaedros compartilhando arestas. ....</b>	<b>239</b>
5.44 HONESSITA $(\text{Ni}_{1-x}\text{Fe}^{3+}_x)(\text{SO}_4)_{x/2}(\text{OH})_2 \cdot n\text{H}_2\text{O} (x < 0,5; n < 3/x/2) - 7.\text{DD}.25$ .....	239
<b>7.DE. Sulfatos (Teluratos) com cátions de raios atômicos médios; não classificados. ....</b>	<b>241</b>
5.45 BRUMADOÍTA $\text{Cu}_3(\text{Te}^{6+}\text{O}_4)(\text{OH})_4 \cdot 5\text{H}_2\text{O} - 7.\text{DE}.80$ .....	241
<b>7.DF. Sulfatos com cátions de raios atômicos médios e grandes.....</b>	<b>242</b>
5.46 CAINITA $\text{KMg}(\text{SO}_4)\text{Cl} \cdot 3\text{H}_2\text{O} - 7.\text{DF}.10$ .....	242
5.47 METAVOLTINA $\text{K}_2\text{Na}_6\text{Fe}^{2+}\text{Fe}^{3+}_6\text{O}_2(\text{SO}_4)_{12} \cdot 18\text{H}_2\text{O} - 7.\text{DF}.35$ .....	244
5.48 DUKEÍTA $\text{Bi}^{3+}_{24}\text{Cr}^{6+}_8\text{O}_{57}(\text{OH})_6 \cdot 3\text{H}_2\text{O} - 7.\text{DF}.80$ .....	245
<b>7.E. Sulfatos com uranila. ....</b>	<b>247</b>
<b>7.EC. Sulfatos com cátions de raios atômicos médios a grandes.....</b>	<b>247</b>
5.49 ZIPPEÍTA $\text{K}_3(\text{UO}_2)_4(\text{SO}_4)_2\text{O}_3(\text{OH}) \cdot 3\text{H}_2\text{O} - 7.\text{EC}.05$ .....	247

The four volumes (in Portuguese) are part of a monumental work that covers practically all the mineral species present in Brazil. The work consists of six volumes and four above have already been published.

The two authors have a vast professional experience.

According to the International Mineralogical Association 4893 mineral species have been validated world wide up to the present date. Of these, 894 species have been registered in Brazil. This monumental work represents the major compilation of minerals present in Brazil and is of extreme scientific importance, not only for Brazil.

Some fifty years ago, and Old American Indian Friend from New Mexico – The Land of Enchantment, told us that there are four things in Nature and the Universe that fascinate mankind with their beauty, radiance and magic. They are the *stars, the flowers, the minerals and the sparkling eyes of beautiful women*. (Cf. L. G. Ionescu, South. Braz. J. Chem., 11(12), 22, 2003).

*(The 4<sup>th</sup> volume of Enciclopédia dos minerais do Brasil. with Carbonates, Sulfates, and Organic Compounds, is a considerable improvement of this monumental work written by Neves and Atencio. We congratulate the authors, and highly recommend this work to all persons interested in the Earth Sciences.)*

L. G. Ionescu. B. S., M. S., Ph. D. (Physical Chemistry/Astrophysics)  
Sarmisegetuza Research Group. Santa Fe, New Mexico, USA.

Enciclopedia dos Minerais do Brasil, Volume 4, Editora da ULBRA, 2015, 392 pp.

De acordo com a *International Mineralogical Association* 4893 espécies minerais foram validadas até o presente à nível global. Destas, 894 espécies foram registradas no Brasil. A realização desta obra monumental que representa a maior compilação de minerais existentes no Brasil é de máxima importância para a ciência no nosso País.

Os dois autores possuem ampla experiência profissional.

Cinqüenta anos atrás, um Velho Índio Americano, nosso amigo do Novo México – Terra do Encanto, dizia que tem quatro coisas que são verdadeiramente bonitas e extraordinárias na Natureza e no Universo, que fascinam e encantam com a sua beleza, brilho e magia. São *as estrelas, as flores, os minerais e os olhos cintilantes de mulheres bonitas*. (Cf. L.G. Ionescu, *South. Braz. J. Chem.*, 11(12), 22, 2003).

A presente obra com seu alto nível científico e altíssima qualidade gráfica nos traz uma das partes mais fascinantes do Universo Brasileiro.

O quarto volume da *Enciclopédia dos minerais do Brasil. com Carbonatos, Sulfatos e Combinações Orgânicas*, é um considerável incremento nesta monumental obra que está sendo escrita por Neves e Atencio. Congratulamo-nos com os autores e recomendamos este trabalho a todos aqueles que se interessam pelas Ciências da Terra.

*Lavinell G. Ionescu, A.A., B.S., M.S., Ph.D. (Físico-Química/Astrofísica)*

## AUTHOR INDEX / ÍNDICE DE AUTORES

Aboaba, S. A.	23, 83
Adeeko, O.	23
Adegbe, A. A.	73
Bolaji, O. S.	37
Dada, A. O.	61
Dalloul, H. M.	49
El-nwairy, K.	49
Ekundayo, O.	83
Flamini, G.	23
Ionescu, L. G.	02
Kath, C.	02
Larayetan, R. A.	73
Latona, D. F.	61
Neves, P. C. P. das	02
Ogunmola, O. O.	37, 91
Owonikoko, A. D.	73
Samaha, A. A.	49
Shorafa, A. Z.	49
Sodamade, A.	37
Ugah, T. C.	83

*SCIENCO*  
**SOUTHERN BRAZILIAN JOURNAL  
OF CHEMISTRY**

ISSN 0104-5431

The **SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY – *SCIENCO*** (**SOUTH. BRAZ. J. CHEM.**) publishes original research articles in chemistry and related interdisciplinary areas and is intended to fill a gap in terms of scientific information for Southern Brazil.

Occasionally the journal will include review papers and articles dealing with chemical education and philosophy and history of science. It will be published mainly in English, with abstracts in Portuguese and only occasional papers in other languages. At the present there are no page charges and the authors will receive twenty five reprints of their papers free of charge.

We have set high standards for the articles to be published by ensuring strong but fair refereeing by at least two reviewers. We hope that this journal will provide a forum for dissemination of high quality research in chemistry and related areas and are open to any questions and suggestions.

The Editor

**SUBSCRIPTION INFORMATION**

Brazil and Latin America:

US\$ 70.00 per issue,

Other Countries: US\$ 100.00 per issue,

including air mail delivery. Persons or institutions outside Brazil should send subscription fee payable to Dr. L. G. Ionescu,

c/o SBJC, 8532 Howard Circle, Huntington Beach, California

USA 92647

**MAILING ADDRESS**

***SOUTHERN BRAZILIAN JOURNAL OF CHEMISTRY***

Lavinel G. Ionescu, B.S., M.S., Ph.D., Editor

C.P. 15032, Agronomia

Porto Alegre, RS, BRASIL 91501-970

Tel. 055 51 3485-1820 / 051 55 3485-1711 / 055 51 995-26616 / 055 51997-64159



E-Mail: [lavinel@gmail.com](mailto:lavinel@gmail.com)

VISIT OUR SITE: [www.sbjchem.tk](http://www.sbjchem.tk)

**FINANCIAL SUPPORT**

**SARMISEGETUSA RESEARCH GROUP**

**SANTA FE, NEW MEXICO, U.S.A.**



Endless Column, 1937, cast iron  
CONSTANTIN BRĂNCUȘI